

522984



Sandia National Laboratories

Operated for the U.S. Department of Energy by Sandia Corporation

Carlsbad, New Mexico 88220

date: July 26, 2002

to: Laurence H. (Larry) Brush
Geochemistry Team Lead
Org. 6822, MS-1395
Carlsbad Programs Group

from: Emily R. Giambalvo
Performance Assessment
Org. 6821, MS-1395
Carlsbad Programs Group

subject: Recommended Parameter Values for Modeling An(IV) Solubility in WIPP Brines

1. Introduction

Parameter values for modeling speciation and solubility of Th(IV) from the recent literature (Felmy and Rai, 1999) offer the opportunity to improve the speciation and solubility model for actinides in the +IV oxidation state (An(IV)) that is implemented in the Fracture Matrix Transport (FMT) geochemical database. The newly available parameters are the ternary ion interaction parameters (θ and ψ) describing the interaction of $\text{Th}(\text{CO}_3)_5^{6-}$ with Cl^- and Na^+ .

This memo describes the implementation of the An(IV) speciation and solubility model in the version of the FMT database used for compliance baseline performance assessment (PA) calculations (baseline database)*, and recommends modifications to the two parameters listed above. It also recommends modification of the value used for the normalized chemical potential (μ^0/RT , where R is the ideal gas constant and T is temperature in Kelvin) of $\text{Th}(\text{OH})_{4(\text{aq})}$.

2. An(IV) Model in the Baseline Database

The An(IV) speciation and solubility model in the current baseline database is that described by Novak (1996; 1997). Parameters are derived from Th solubility data and electromotive force measurements. The WIPP actinide solubility model assumes that Th(IV) solubilities predicted by the An(IV) model are representative of Pu(IV) and U(IV) solubility in addition to Th(IV) solubility (DOE, 1996). This assumption is conservative, i.e. Th(IV) is more soluble than Pu(IV) and U(IV) (e.g., Neck and Kim, 2001; Wall et al., 2002). The parameters used in the baseline An(IV) model are derived as follows:

1. Hydrolysis species (except $\text{Th}(\text{OH})_{4(\text{aq})}$) and polynuclear complexes of Th are neglected. The solubility product constant (K_{sp}) for $\text{ThO}_{2(\text{am})}$ was calculated by Felmy et al. (1991) from Th solubility data collected in dilute solution at $\text{pH} < 7$ (Ryan and Rai, 1987). Felmy et al. (1991)

* The baseline database is identical to the database used in the Performance Assessment Verification Test (PAVT), and is the most recent documented version of the database (Novak, 1997). It is stored in the CHEMDAT file FMT_970407.CHEMDAT in Sandia National Laboratories' Configuration Management System on the WIPP Alpha Cluster in the library WP\$NONPA_CMSROOT:[FMT].

Information Only
Exceptional Service in the National Interest

WIPP:1.3.5.1.2.1:PA:QA:FMT Database for CRA

neglected all hydrolysis species in their calculation of K_{sp} . They successfully simulated Th solubility in NaCl solutions up to 3 mol/L using their calculated K_{sp} and Pitzer parameters describing the interaction of Th^{4+} and Cl^- from Pitzer and Mayorga (1973).

In order to model $\text{ThO}_{2(\text{am})}$ solubility at $\text{pH} > 7$, Novak (1996) added $\text{Th}(\text{OH})_{4(\text{aq})}$ to the model. In the pH range 7-14, Th solubility is pH-independent and $\text{Th}(\text{OH})_{4(\text{aq})}$ is the dominant Th species in solution (in the absence of carbonate) (Ryan and Rai, 1987; Neck and Kim, 2001). Novak (1996) cites only "new work by PNL [Pacific Northwest National Laboratory]" as the source for μ^0/RT of $\text{Th}(\text{OH})_{4(\text{aq})}$. I was unable to discover the data set from which this value is derived.

2. Th-carbonate complexes include $\text{Th}(\text{CO}_3)_5^{6-}$ and $\text{Th}(\text{OH})_3\text{CO}_3^-$. Novak (1996) determined μ^0/RT values for these species by comparing $\text{ThO}_{2(\text{am})}$ solubility in the presence of carbonate (Felmy et al., 1997) to $\text{ThO}_{2(\text{am})}$ solubility in the absence of carbonate (Felmy et al., 1991). Felmy et al. (1997) determined the solubility of $\text{ThO}_{2(\text{am})}$ in the presence of carbonate from solubility experiments run in dilute to moderately concentrated (< 2 mol/L) NaHCO_3 , Na_2CO_3 , and $\text{NaClO}_4/\text{CO}_{2(\text{g})}$ solutions. Pitzer parameters describing interaction of $\text{Th}(\text{CO}_3)_5^{6-}$ with Na^+ ($\beta^{(0)}$, $\beta^{(1)}$, and C^ϕ) and with ClO_4^- (θ) are from Felmy et al. (1997). Pitzer parameters describing the ternary interaction of $\text{Th}(\text{CO}_3)_5^{6-}$, Cl^- , and Na^+ (θ , ψ) result from A. Felmy's preliminary analysis of $\text{ThO}_{2(\text{am})}$ solubility in mixed $\text{NaCl}/\text{Na}_2\text{CO}_3$ solutions (Novak, 1997).
3. Values of μ^0/RT for Th-sulfate complexes, related Pitzer parameters, and K_{sp} values for Th-sulfate solid phases were derived from various data sets by Felmy and Rai (1992). Like Felmy et al. (1991), Felmy and Rai (1992) used the Pitzer and Mayorga (1973) Th^{4+} - Cl^- interaction parameters in their calculations.
4. Th-chloride complexes are not explicitly modeled. Instead they are accounted for through appropriate Pitzer ion interaction parameters. The Th^{4+} - Cl^- ion interaction parameters included in the baseline database were determined from electromotive force measurements in mixed HCl/ThCl_4 solutions (Roy et al., 1992). Although this set of Pitzer parameters is not identical to that used by Felmy et al. (1991) and Felmy and Rai (1992), Roy et al. (1992) used it to successfully simulate the systems studied by Felmy et al. (1991) and Felmy and Rai (1992) without altering μ^0/RT values or K_{sp} s.
5. Pitzer parameters describing the ternary interaction of Th^{4+} , H^+ , and Cl^- are from the electromotive force measurements of Roy et al. (1992).
6. Pitzer parameters describing the interaction of Th^{4+} with Na^+ and Mg^{2+} were calculated from $\text{ThO}_{2(\text{am})}$ solubility data collected in NaCl (up to 6 m) and MgCl_2 (up to 3 m) brines (Rai et al., 1997). Rai et al. (1997) used the K_{sp} determined by Felmy et al. (1991) and the Roy et al. (1992) Th^{4+} - Cl^- Pitzer parameters in their calculations.

Tables 1-4 list the parameter values used in the baseline An(IV) model.

3. Recommended An(IV) Model

The recommended An(IV) model improves on the baseline An(IV) model in two ways:

Information Only

1. Pitzer parameters describing the ternary interaction of $\text{Th}(\text{CO}_3)_5^{6-}$, Cl^- , and Na^+ (θ , ψ) come from a peer-reviewed article (Felmy and Rai, 1999), rather than an informal communication (e-mail from A. Felmy cited in Novak, 1997). They are derived from $\text{ThO}_{2(\text{am})}$ solubility data collected in mixed $\text{Na}_2\text{CO}_3/\text{NaCl}$ solutions (2.33 and 4.67 mol/kg NaCl).
2. The μ^0/RT value for $\text{Th}(\text{OH})_{4(\text{aq})}$ is derived from a traceable data set (Ryan and Rai, 1987) as follows. From the low pH (<7) data of Ryan and Rai (1987) Felmy et al. (1991) calculated $\log K_{\text{sp}} = -45.5$ for the dissolution reaction:



In the pH range where $\text{Th}(\text{OH})_{4(\text{aq})}$ is the dominant Th species in solution (pH = 7-14), Ryan and Rai's (1987) data yield a Th solubility of $10^{-8.8}$ mol/L, or:



Differencing these equations yields the complexation constant for $\text{Th}(\text{OH})_{4(\text{aq})}$, which is easily converted to μ^0/RT :



This calculation assumes that all Pitzer parameters related to $\text{Th}(\text{OH})_{4(\text{aq})}$ are zero (i.e., the activity of $\text{Th}(\text{OH})_{4(\text{aq})}$ is independent of ionic strength). The complexation constant obtained in this way is similar to that used in the baseline An(IV) model and to that obtained from a similar analysis of the solubility data presented by Felmy et al. (1991) ($\log K = 36.9$ and 37.0 , respectively). The $\log K$ calculated from either the Ryan and Rai (1987) or the Felmy et al. (1991) data set results in a conservative prediction of Th solubility at neutral and alkaline pH, because, in both experiments, measured solubilities in the pH range 7-14 were at the detection limit of the measurement technique.

Tables 1-4 list the recommended parameter values and compare them to the values used in the baseline version of the database.

4. References

- DOE, 1996. Appendix SOTERM in *Title 40 CFR Part 191 Compliance Certification Application for the Waste Isolation Pilot Plant*, US Department of Energy, Carlsbad, NM.
- Felmy, A.R., D. Rai and M.J. Mason, 1991. The solubility of hydrous thorium(IV) oxide in chloride media: Development of an aqueous ion-interaction model, *Radiochimica Acta*, **55**:177-185. WPO 40225.
- Felmy, A.R. and D. Rai, 1992. An aqueous thermodynamic model for a high valence 4:2 electrolyte $\text{Th}^{4+}-\text{SO}_4^{2-}$ in the system $\text{Na}^+-\text{K}^+-\text{Li}^+-\text{NH}_4^+-\text{Th}^{4+}-\text{SO}_4^{2-}-\text{HSO}_4^--\text{H}_2\text{O}$ to high concentration, *Journal of Solution Chemistry*, **21**:407-423.

- Felmy, A.R., D. Rai, M.S. Sterner, M.J. Mason, N.J. Hess and S.D. Conradson, 1997. Thermodynamic models for highly charged aqueous species: Solubility of Th(IV) hydrous oxide in concentrated NaHCO₃ and Na₂CO₃ solutions, *Journal of Solution Chemistry*, **26**:233-248.
- Felmy, A.R. and D. Rai, 1999. Application of Pitzer's equations for modeling the aqueous thermodynamics of actinide species in natural waters: A review, *Journal of Solution Chemistry*, **28**:533-553. SAND99-0340J
- Fuger, J. and F.L. Oetting, 1976. *The Chemical Thermodynamics of Actinide Elements and Compounds. Part 2. The Actinide Aqueous Ions.*, Int. Atomic Energy Agency, Vienna, Austria.
- Neck, V. and J.I. Kim, 2001. Solubility and hydrolysis of tetravalent actinides, *Radiochimica Acta*, **89**:1-16.
- Novak, C.F., 1996. "Preliminary inorganic model for thorium solubility in WIPP brines, in database file HMW_3Th5_960119.CHEMDAT," Memo to E. J. Nowak, 19 January 1996, Albuquerque, NM: Sandia National Laboratories. WPO #30930
- Novak, C.F., 1997. "Calculation of actinide solubilities in WIPP SPC and ERDA-6 brines under MgO backfill scenarios containing either nesquehonite or hydromagnesite as the Mg-CO₃ solubility-limiting phase," Memo to R. V. Bynum, April 21, 1997, Albuquerque, NM: Sandia National Laboratories. WPO 46124
- Pitzer, K.S. and G. Mayorga, 1973. Thermodynamics of electrolytes. II. Activity and osmotic coefficients for strong electrolytes with one or both ions univalent, *The Journal of Physical Chemistry*, **77**:2300-2308.
- Rai, D., A.R. Felmy, M.S. Sterner, D.A. Moore, M.J. Mason and C.F. Novak, 1997. The solubility of Th(IV) and U(IV) hydrous oxides in concentrated NaCl and MgCl₂ solutions, *Radiochimica Acta*, **79**:239-247. SAND96-2909J, WPO 51469.
- Roy, R.N., K.M. Vogel, C.E. Good, W.B. Davis, L.N. Roy, D.A. Johnson, A.R. Felmy and K.S. Pitzer, 1992. Activity coefficients in electrolyte mixtures: HCl + ThCl₄ + H₂O for 5-55 °C, *J. Phys. Chem.*, **96**:11065-11072.
- Ryan, J.L. and D. Rai, 1987. Thorium(IV) Hydrous Oxide Solubility, *Inorganic Chemistry*, **26**:4140-4142.
- Wall, N.A., E.R. Giambalvo, L.H. Brush and D.E. Wall, 2002. "The Use of Oxidation-State Analogs for WIPP Actinide Chemistry," 223rd American Chemical Society National Meeting, Orlando, FL, 2 pp. SAND2002-0698P.

ERG:6821:erg/(2002-1007, Rev. A)

Copy to:

MS-1395, M.K. Knowles [Dept. 6821]

MS-1395, D.E. Wall [Dept. 6822]

MS-1395, N.A. Wall [Dept. 6822]

Information Only

Laurence H. (Larry) Brush
Org. 6822, MS-1395

- 5 -

July 26, 2002

MS-1395, Y. Xiong [Dept. 6822]
MS-1395, Y. Wang [Dept. 6823]
MS-0779, R.C. Moore [Dept. 6849]
MS-1395, E.R. Giambalvo [Dept. 6821]
MS-1395, Day File [Dept. 6821]

Information Only

Table 1. Normalized Chemical Potential (μ^0/RT)

Species	Current Baseline		Recommended	
	μ^0/RT	Source	μ^0/RT	Source
Th ⁴⁺	-284.227	Fuger and Oetting (1976)		<i>same</i>
Th(CO ₃) ₅ ⁶⁻	-1411.378	Novak (1996), based on Felmy et al. (1997)		<i>same</i>
Th(OH) ₃ (CO ₃) ⁻	-775.627	Novak (1996), based on Felmy et al. (1997)		<i>same</i>
Th(OH) _{4(aq)}	-622.84	Novak (1996)	-622.47	Present work, based on Ryan and Rai (1987)
Th(SO ₄) _{2(aq)}	-911.69	Felmy and Rai (1992)		<i>same</i>
Th(SO ₄) ₃ ²⁻	-1214	Felmy and Rai (1992)		<i>same</i>
ThO _{2(am)}	-451.408	Felmy et al. (1991)		<i>same</i>
Th(SO ₄) ₂ ·9H ₂ O _(s)	-1775.9	Felmy and Rai (1992)		<i>same</i>
Th(SO ₄) ₂ ·8H ₂ O _(s)	-1680	Felmy and Rai (1992)		<i>same</i>
Th(SO ₄) ₂ ·Na ₂ SO ₄ ·6H ₂ O _(s) ^a	-2011.29	Felmy and Rai (1992)		<i>same</i>
Th(SO ₄) ₂ ·K ₂ SO ₄ ·4H ₂ O _(s) ^a	-1837.57	Felmy and Rai (1992)		<i>same</i>
Th(SO ₄) ₂ ·2K ₂ SO ₄ ·2H ₂ O _(s) ^a	-2181.81	Felmy and Rai (1992)		<i>same</i>
Th(SO ₄) ₂ ·3.5K ₂ SO _{4(s)} ^a	-2790.83	Felmy and Rai (1992)		<i>same</i>

^a μ^0/RT value is at 16°C. Unless noted, all values in the FMT geochemical database are at 25°C.

Table 2. Binary Pitzer Parameters ($\beta^{(0)}$, $\beta^{(1)}$, $\beta^{(2)}$, C^ϕ)^a

Cation	Anion	$\beta^{(0)}$	$\beta^{(1)}$	$\beta^{(2)}$	C^ϕ	Source
Na ⁺	Th(SO ₄) ₃ ²⁻	0.12	0	0	0	Felmy and Rai (1992)
Na ⁺	Th(OH) ₃ (CO ₃) ⁻	0	0	0	0	Felmy et al. (1997)
Na ⁺	Th(CO ₃) ₅ ⁶⁻	1.31	30	0	0	Felmy et al. (1997)
K ⁺	Th(SO ₄) ₃ ²⁻	0.9	0	0	0	Felmy and Rai (1992)
H ⁺	Th(SO ₄) ₃ ²⁻	0.84	0	0	0	Felmy and Rai (1992)
Th ⁴⁺	Cl ⁻	1.092	13.7	-160	-0.112	Roy et al. (1992)
Th ⁴⁺	SO ₄ ²⁻	1.56	0	0	0	Felmy and Rai (1992)
Th ⁴⁺	HSO ₄ ⁻	1.44	0	0	0	Felmy and Rai (1992)
Th ⁴⁺	ClO ₄ ⁻	1.19	27.3	0	-0.057	Novak (1996) ^b

^aRecommended values are identical to those used in the current baseline model.

^bNovak (1996) references "new work, SNL/FSU" for these values. Although the exact source is unclear, I did not re-evaluate Th⁴⁺-ClO₄⁻ parameters, because ClO₄⁻ is not a component in simulations of WIPP brines.

Handwritten notes and signatures in the bottom right corner.

Table 3. Neutral-Ion Interaction Parameter (λ)^a

Neutral	Ion	λ	Source
Th(SO ₄) _{2(aq)}	Cl ⁻	0.29	Felmy and Rai (1992)
Th(SO ₄) _{2(aq)}	HSO ₄ ⁻	0.68	Felmy and Rai (1992)

^aRecommended values are identical to those used in the current baseline model.

Table 4. Ternary Interaction Parameters (θ , ψ)

ion <i>i</i>	ion <i>j</i>	ion <i>k</i>	Current Baseline			Recommended		
			θ_{ij}	ψ_{ijk}	Source	θ_{ij}	ψ_{ijk}	Source
Na ⁺	Th ⁴⁺	Cl ⁻	0.42	0.21	Rai et al. (1997)			<i>same</i>
Mg ²⁺	Th ⁴⁺	Cl ⁻	0.60	0.21	Rai et al. (1997)			<i>same</i>
H ⁺	Th ⁴⁺	Cl ⁻	0.60	0.37	Roy et al. (1992)			<i>same</i>
Th(CO ₃) ₅ ⁶⁻	Cl ⁻	Na ⁺	2.0	-0.08	Novak (1997)	1.8	0.3	Felmy and Rai (1999)
Th(CO ₃) ₅ ⁶⁻	ClO ₄ ⁻	Na ⁺	5.5	0	Felmy et al. (1997)			<i>same</i>

Information Only

-9-