



GLOSSARY OF TERMS

1
2
3 **40 CFR Part 191.** *Environmental Radiation Protection Standards for Management and*
4 *Disposal of Spent Nuclear Fuel, High-Level and Transuranic Radioactive Wastes.* This
5 regulation sets environmental radiation protection standards for management (Subpart
6 A) and disposal (Subparts B and C) of spent nuclear fuel and high-level and transuranic
7 radioactive wastes.

8
9 **40 CFR Part 194.** *This regulation provides EPA's criteria for the certification and*
10 *recertification of the Waste Isolation Pilot Plant's compliance with 40 CFR Part 191*
11 *Disposal Regulations.*

12
13 **40 CFR Part 261.** *Identification and Listing of Hazardous Waste.* This part identifies those
14 solid wastes which are subject to regulation as hazardous wastes under Parts 262–265,
15 268, 270, 271, and 124 of Title 40 of the Code of Federal Regulations.

16
17 **40 CFR Part 264.** *Standards for Owners and Operators of Hazardous Waste Treatment,*
18 *Storage, and Disposal Facilities.* This subpart establishes minimum national standards
19 which define the acceptable management of hazardous waste.

20
21 **40 CFR Part 264.** Subpart G. This subpart of 40 CFR Part 264 defines closure and post-
22 closure requirements pertaining to hazardous waste management units.

23
24 **40 CFR Part 264.** Subpart X. This subpart specifies requirements that apply to owners and
25 operators of facilities that treat, store, or dispose of hazardous waste in miscellaneous
26 hazardous waste management units.

27
28 **40 CFR Part 268.** This regulation restricts the land disposal of hazardous waste and specifies
29 treatment standards and/or treatment technologies that must be met or applied before
30 hazardous wastes may be land disposed. Section 268.6 provides for petitioning to allow
31 land disposal of untreated hazardous waste if it can be demonstrated to a reasonable
32 degree of certainty that there will be no migration of hazardous constituents from the
33 disposal unit for as long as the waste remains hazardous.

34
35 **40 CFR Part 270.** This regulation establishes provisions for the Hazardous Waste Permitting
36 Program under Subtitle C of Resource Conservation and Recovery Act. This regulation
37 and the associated State of New Mexico regulation require the permitting of the WIPP as
38 a hazardous waste facility.

39
40 **abscissa.** The horizontal coordinate (x coordinate) of a point plotted in Cartesian (x, y)
41 coordinates.



1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44

absolute temperature. Temperature measured in degrees Celsius from absolute zero, -273.18°C. Absolute temperatures are given either as “degrees absolute” (for example, 150° A) or as degrees kelvin (for example, 150 K).

absorb. The process by which a fluid penetrates or is drawn into and held in the inner structure of another (solid or liquid) material. The word also appears in contexts such as absorbed energy and absorbed radiation dose, in which case it refers to internal energy imparted to the absorbing material. Absorbed radiation dose refers specifically to the energy imparted by ionizing radiation per unit mass of irradiated material at a particular location.

accessible environment. “(1) [T]he atmosphere, (2) land surfaces, (3) surface waters, (4) oceans, and (5) all of the lithosphere that is beyond the controlled area.” (40 CFR § 191.12)

acid rain. Any form of precipitation having a pH of 5.6 or less. Major components contributing to the acid rain are sulfur dioxide and oxides of nitrogen.

actinide intrinsic colloids. Macromolecules or aggregations of actinide compounds which form by exsolution of dissolved actinide compounds and remain in suspension thereby acting as colloids.

actinide source term. Modeling terminology denoting the fraction of total actinide inventory of a disposal room or the repository that can be mobilized for transport.

active institutional control. (1) Controlling access to a disposal site by any means other than passive institutional controls, (2) performing maintenance operations or remedial actions at a site, (3) controlling or cleaning up releases from a site, or (4) monitoring parameters related to disposal system performance. (40 CFR § 191.12)

activity. A measure of the rate at which a material emits nuclear radiation, usually given in terms of the number of nuclear disintegrations occurring in a given length of time. The unit of activity used in this document is the curie (Ci).

adsorption. (1) The net accumulation of matter at the interface between a solid phase and an aqueous solution through the formation of two-dimensional (that is, noncrystallographic) molecular structures. The accumulated matter is the *adsorbate*. The solid surface is the *adsorbent*. (2) Adherence of gas molecules to the surface of solids in which they are in contact.

advection. The transport of matter (for example, a trace contaminant) by motion of the medium in which it is contained. The most common transport medium is the flow of a fluid.



- 1 **aeolian.** Also spelled eolian. Applies to the erosional action of wind, and to deposits which
2 are due to the transporting action of wind.
3
- 4 **aerobic degradation.** The microbial degradation of organic (that is, cellulosic, rubber,
5 plastic, etc.) wastes using oxygen (O₂) as the electron acceptor and producing carbon
6 dioxide (CO₂) and water (H₂O).
7
- 8 **aeromagnetic survey.** Refers to the geomagnetic observations using a magnetometer from an
9 airborne craft.
10
- 11 **air lock.** An intermediate chamber between zones of different static pressure.
12
- 13 **alluvial.** Pertaining to or composed of alluvium, or deposited by a stream or running water,
14 for example, an alluvial clay or an alluvial divide.
15
- 16 **algal features.** Features created by the secretion of calcium carbonate from algae.
17
- 18 **alkalophilic bacteria.** Bacteria that can grow and reproduce well in an alkaline (high pH)
19 environment.
20
- 21 **alpha emitter.** A radioactive isotope that decays through the emission of alpha particles.
22
- 23 **alpha particle.** (α -particle). A positively charged particle emitted in the radioactive decay of
24 certain nuclides. Made up of two protons and two neutrons bound together, it is
25 identical to the nucleus of a helium atom. It is the least penetrating of the three common
26 types of ionizing radiation—alpha, beta, and gamma radiation.
27
- 28 **alternative conceptual models.** Alternative sets of assumptions that describe the same
29 system for the same purpose and are consistent with the existing information.
30
- 31 **anaerobic.** The microbial degradation of organic (that is, cellulosic, rubber, plastic, etc.)
32 wastes in the absence of oxygen (O₂), usually consuming nitrate (NO₃) as the electron
33 acceptor and producing carbon dioxide (CO₂), nitrous oxide (N₂O), and nitrogen (N₂).
34
- 35 **anhydrite.** A mineral consisting of anhydrous calcium sulfate (CaSO₄). It is gypsum without
36 water and is denser, harder, and less soluble.
37
- 38 **annual committed effective dose.** The committed effective dose resulting from a one-year
39 intake of radionuclides released plus the annual effective dose caused by direct radiation
40 from facilities or activities subject to Subparts B and C of 40 CFR Part 191. (40 CFR
41 § 191.12)
42



- 1 **annulus.** The area between the borehole wall and the outside of the drill pipe in rotary
2 drilling operations. Rotary drilling is used for scientific investigation studies, such as
3 the WIPP project, and the exploration for natural resources (that is, oil and gas).
4
- 5 **anoxic.** Describes a process that occurs in the absence of oxygen (O₂). Anoxic corrosion of
6 steels, iron-base alloys, and other metal wastes use water (H₂O) or hydrogen sulfide
7 (H₂S) and produces hydrogen gas (H₂).
8
- 9 **anthropogenic.** Resulting from the influence of human beings on nature (that is, acid rain,
10 greenhouse effect, man-made organic material).
11
- 12 **anticline.** A fold of rocks whose core contains the stratigraphically older rocks; it is
13 generally convex upward.
14
- 15 **aquifer.** An underground geological formation or part of a formation that is capable of
16 yielding a significant amount of water to a well or spring. (40 CFR § 191.12)
17
- 18 **arenaceous.** Of the texture or character of sand.
19
- 20 **argillaceous rocks.** A group of detrital sedimentary rocks with a grain or particle size less
21 than 1/16 millimeter. Commonly comprised of clay minerals and including shales,
22 mudstones, siltstones, and marls.
23
- 24 **artesian.** Adjective that refers to groundwater confined under pressure so that it will rise in a
25 well. Sometimes the word is used to mean that the water flows out at the surface, but
26 that, strictly speaking, is “flowing artesian.”
27
- 28 **assurance requirements.** Qualitative requirements in 40 CFR § 191.14 that specify actions
29 and procedures to increase confidence that the probabilistic release events in the
30 containment requirements (40 CFR § 191.13) will be met.
31
- 32 **Atoka.** The Early-Middle Pennsylvanian rocks, assigned to the Atokan or Derryan Stage,
33 consist of dark-colored sandstones, shales and limestones, which attain a maximum
34 thickness of about 1,000 feet.
35
- 36 **atomic weights.** Atomic weight is the total mass of an atom and is almost, but not exactly,
37 the sum of the masses of its constituent protons, neutrons and electrons.
38
- 39 **attenuation.** (1) A reduction in the amplitude or energy of a signal, such as might be
40 produced by passage through a filter. (2) A reduction in the amplitude of seismic waves,
41 as produced by divergence, reflection and scattering, and absorption. (3) That portion of
42 the decrease in seismic or sonar signal strength with distance that is not dependent on
43 geometrical divergence, but on the physical characteristics of the transmitting medium.
44



1 **authigenic.** Formed or generated in place of rock constituents and minerals that have not
2 been transported or that crystallized locally at the spot where they are now found, and of
3 minerals that came into existence at the same time as, or subsequently to, the formation
4 of the rock of which they constitute a part. The term often refers to a mineral (such as
5 quartz or feldspar) formed after deposition of the original sediment.
6

7 **backfill.** Material placed around or above the waste containers, filling the open space in the
8 disposal room.
9

10 **background (radiation).** Radiation in the human environment from naturally occurring
11 elements, fallout, and from cosmic radiation.
12

13 **bacterioruberin.** A carotenoid pigment excreted during growth and multiplication of various
14 strains of bacteria that may be involved in waste decomposition.
15

16 **barrier.** “[A]ny material or structure that prevents or substantially delays movement of water
17 and/or radionuclides toward the accessible environment. For example, a barrier may be
18 a geologic structure, a canister, a waste form with physical and chemical characteristics
19 that significantly decrease the mobility of radionuclides, or a material placed over and
20 around waste, provided that the material or structure substantially delays movement of
21 water or radionuclides.” (40 CFR § 191.12) Barriers also prevent or delay the
22 movement of hazardous constituents.
23

24 **Baseline Inventory Report.** Also known as the Transuranic Waste Baseline Inventory
25 Report. This document provides the total DOE transuranic waste inventory. The
26 Baseline Inventory Report establishes a methodology for grouping wastes of similar
27 physical and chemical properties from across the DOE transuranic waste system into a
28 series of waste profiles.
29

30 **bedding plane.** A flat or near flat bedding surface that visibly separates the individual layers,
31 beds, or strata in sedimentary or stratified rocks from the preceding or following layers.
32

33 **Bell Canyon Formation.** A sequence of rock strata that form the uppermost formation of the
34 Delaware Mountain Group (Early Permian). It is immediately below the Castile
35 Formation at about 4,000 feet below the surface at the WIPP. May contain some oil and
36 gas.
37

38 **bentonite.** A commercial term applied to expansive clay materials containing
39 montmorillonite (smectite) as the essential mineral.
40

41 **beta particle.** (β -particle). A negatively charged particle emitted in the radioactive decay of
42 certain radioactive elements (nuclides). A beta particle is identical to an electron. It is
43 not a strong penetrating radiation and can be stopped by an inch of wood or a thin sheet
44 of aluminum.



- 1 **biodegradation.** The process of consumption by microbial substances—usually organic
2 materials such as cellulosics.
3
- 4 **bioelectric activity.** The electric phenomena associated with organic matter (plants and
5 animals).
6
- 7 **biofilm ecosystem.** An ecosystem composed of substrates and nutrients developed on
8 surfaces containing one or more layers of cells with extra cellular polymeric material.
9
- 10 **biological half-life.** The time required for an organism to eliminate half the amount of a
11 radionuclide ingested or inhaled. See definition for half-life.
12
- 13 **blowout.** An unexpected volume of gas or fluid released from subsurface pressurized zones
14 that flows upwards to the land surface and into the atmosphere during the drilling of a
15 borehole.
16
- 17 **Boltzmann Constant.** The ideal gas constant per molecule is the ratio of the molar gas
18 constant to the number of molecules per gram of a substance. Boltzman's Constant is
19 defined as having a value of approximately 1.380×10^{-23} Joules/Molecule· K.
20
- 21 **borehole.** (1) A hole drilled from the surface for purposes of geologic or hydrologic testing,
22 injection, or exploration and subsequent production of resources; sometimes referred to
23 as a drillhole or well bore. (2) A man-made hole in the wall, floor, or ceiling of a
24 subsurface room used for verifying geology, making observations, or emplacing
25 canisters of remote-handled transuranic waste.
26
- 27 **brackish.** Water with between 1 and 10 parts per thousand salinity. This places the saline
28 content of brackish waters between that of streams and sea water. See saline and brine.
29
- 30 **BRAGFLO.** The name of the computer model the DOE uses to simulate brine and gas flow
31 in the disposal system.
32
- 33 **breccia.** Fragmented rock whose components are angular and cemented together by a mineral
34 cement or a fine-grained matrix. May be sedimentary (for example, carbonate or
35 evaporite karst features) or formed by crushing or grinding along faults. There are
36 friction or fault breccias, talus breccias, eruptive breccias, and large-scale collapse sinks
37 (for example, breccia pipes). Breccia pipes are vertical or near vertical cylindrical
38 features filled with collapse debris.
39
- 40 **brine.** Water with greater than 100 parts per thousand salinity. Brines are often found in deep
41 sedimentary basins and contain calcium (Ca), sodium (Na), potassium (K), chloride (Cl),
42 sulfate (SO₄), and minor amounts of other elements.
43
- 44 **brine pocket.** See brine reservoir.



1 **brine reservoir.** A volume of brine of limited extent trapped within fractures and/or
2 intergranular pore spaces of a host rock and usually pressurized relative to normal
3 formation fluids. Such pockets may exist under various conditions of stress and solute
4 concentration. Pressurized brine pockets have been observed in the Castile Formation in
5 the Delaware Basin.
6

7 **Brushy Canyon Formation.** The basal unit of the Guadalupian Series or Delaware Mountain
8 Group within the Delaware Basin. The formation is dominated by submarine channel
9 sandstones with minor limestones and shales, and its structural features are indicative of
10 deposition in agitated water. This formation terminates northward against the Bone
11 Spring flexure at the basin margin.
12

13 **Bureau of Land Management.** An agency of the federal government, within the Department
14 of the Interior, which among other land management duties, reviews environmental
15 assessments, environmental impact statements, land acquisition procedures, and any
16 other plans to site waste management facilities on federal lands over which it has
17 jurisdiction.
18

19 **calcareous remnant.** Commonly used with reference to organisms that use calcium
20 compounds to form endo- or exoskeletal structures. The calcareous remnant is that
21 which remains after the soft tissues have disappeared.
22

23 **calcic.** Said of minerals and igneous rocks containing a relatively high proportion of calcium;
24 the proportion required to warrant use of the term depends on circumstances.
25

26 **calcite.** A crystalline mineral composed of calcium carbonate (CaCO_3).
27

28 **calibrate.** To vary parameters of an applied computational model within a reasonable range
29 until differences between observed data and computed values are minimized.
30

31 **caliche.** A calcareous material commonly found in layers on or within the surface of stony
32 soils of arid or semiarid regions. It occurs in the form of gravels, sands, silts, and clays
33 cemented together by calcium carbonate or as crusts at the surface of the soil.
34

35 **Cambrian.** The oldest period of the Paleozoic Era, and the corresponding system of rocks. It
36 is subsequent to the Precambrian and followed by the Ordovician Periods, between
37 approximately 570 and 505 million years ago.
38

39 **canister.** As used in this document, a container, usually cylindrical, for remotely handled
40 transuranic waste. The waste will remain in this canister during and after burial. A
41 canister affords physical containment but not shielding; shielding is provided during
42 shipment by a cask.
43



1 **capillary pressure.** The difference in pressure across the interface between two immiscible
2 fluid phases jointly occupying the interstices of a rock. It is due to the tension of the
3 interfacial surface, and its value depends on the curvature of that surface.
4

5 **Capitan Limestone.** Also referred to as the Capitan Reef and/or Capitan Formation. The
6 Capitan Limestone or Reef is a light colored, fossiliferous and vuggy limestone and
7 breccia which almost completely surrounds the Delaware Basin. The reef was formed in
8 the Early to Mid-Permian Period, between approximately 286 and 260 million years ago.
9 It is a time-stratigraphic equivalent of the Bell Canyon Formation in the Delaware Basin.
10 The reef is a major part of the Guadalupe Mountains from the west to the northwest of
11 the WIPP site, the Apache Mountains and Davis Mountains to the south, and the Glass
12 Mountains to the southeast.
13

14 **Capitan Reef.** See Capitan Limestone.
15

16 **carbonate mudstone.** A nonfissile mud shale consisting mainly of carbonate minerals.
17

18 **carbonates.** Compounds containing the carbonate radical, CO_3^{-2} , primarily applies to rock
19 formations composed of carbonate minerals.
20

21 **Carlsbad potash district.** The area east of Carlsbad formally designated by the U.S.
22 Geological Survey as having potentially economic grades of potash mineralization.
23

24 **carotenoid pigment.** A yellow-to-red color found in microbes and associated biofilms
25 (plants and animals) observed in WIPP microbial gas generation experiments.
26

27 **cask.** A massive shipping container providing shielding for highly radioactive materials and
28 holding one or more canisters.
29

30 **CaSO_4 .** Calcium sulfate.
31

32 **Castile Formation.** A formation of evaporite rocks (mainly anhydrite with a few halite
33 interbeds) of Permian age that immediately underlies the Salado Formation.
34

35 **catalytic surface.** Referring to a substance on a surface that initiates a reaction and enables it
36 to take place at lower temperature or activation energy than in the absence of the
37 catalyst.
38

39 **cation-anion.** A cation is a positively charged ion whereas an anion is a negatively charged
40 ion. A cation in an electrolyzed solution migrates to the cathode (negative terminal); an
41 anion in an electrolyzed solution migrates to the anode (positive terminal).
42



Title 40 CFR Part 191 Compliance Certification Application

1 **cavings.** During rotary drilling, material that erodes from the borehole wall in response to the
2 upward-flowing drilling fluid within the annulus formed by the drill pipe and the
3 borehole wall.
4

5 **cellulosic.** Those materials derived from high polymer plant carbohydrates. Examples are
6 paper, cardboard, wood, cellophane, cloth, etc.
7

8 **cementitious.** A mineral agent, natural or man-made, usually emplaced by hydration or
9 exsolution, that serves to bind together grains of another material, thereby acting as or
10 resembling cement. May also be used to describe a geologic deposit containing a
11 significant amount of cementitious material that was formed by this mechanism.
12

13 **Cenozoic.** An era of geologic time, from the beginning of the Tertiary period to the present
14 (Some authors do not include the Quaternary, considering it a separate era.) It is
15 characterized paleontologically by the evolution and abundance of mammals, advanced
16 mollusks, and birds; paleobotanically, by angiosperms. The Cenozoic is considered to
17 have begun about 66 million years ago.
18

19 **Central Basin Platform.** The geological region known as the High Plains covering an area of
20 several hundreds of square miles separating the Delaware and Midland basins.
21

22 **certification.** Any action taken by the Administrator of the U.S. Environmental Protection
23 Agency under Section 8(d) of the WIPP Land Withdrawal Act.
24

25 **certifier.** In the context of 40 CFR Part 191, the "certifier" is the U.S. Environmental
26 Protection Agency which must certify whether the Department of Energy has
27 demonstrated that the Waste Isolation Pilot Plant is in compliance with the requirements
28 of the standard.
29

30 **chelating agents.** A particular category of complexant that increases solubility by forming a
31 cage around the complexed (or chelated) substance.
32

33 **chemical retardation.** Refers to retardation effects (see Retardation) that are attributable to
34 chemical mechanisms. Chemical reactions with constituents of a rock, precipitation,
35 and sorption of contaminants are examples of chemical retardation.
36

37 **chemical source term.** The fraction of the hazardous constituents inventory that can be
38 mobilized for transport.
39

40 **chemisorption.** The immobilization of a gas, liquid, or material in solution or suspension by
41 the formation of bonds with the surface molecules of a solid material. The strength of
42 these bonds is comparable to that of ordinary chemical bonds.
43



Title 40 CFR Part 191 Compliance Certification Application

1 **Cherry Canyon Formation.** The center sandstone of the Delaware Mountain Group
2 consisting of a fine-grained sandstone to siltstone that is very finely laminated.

3
4 **chert.** A hard, extremely dense or compact, dull to semivitreous, microcrystalline or
5 cryptocrystalline sedimentary rock, consisting dominantly of interlocking crystals of
6 silica (SiO₂) less than about 30 μm in diameter.

7
8 **chronostratigraphic boundaries.** The boundary of a rock stratum that is characterized by
9 having been formed during a specific interval of geologic time.

10
11 **clastic sedimentation.** A sediment formed by the accumulation of fragments derived from
12 preexisting rocks or minerals and transported as separate particles to their places of
13 deposition by purely mechanical agents (such as water, wind, ice, and gravity).

14
15 **clastics.** Pertaining to a rock or sediment composed principally of broken fragments that are
16 derived from preexisting rocks or minerals and that have been transported some distance
17 from their places of origin; also said of the texture of such a rock. The term has been
18 used to indicate a source both within and outside the depositional basin.

19
20 **Clayton Basin.** Northeast of the WIPP site by eight miles. It contains Quaternary alluvium.

21
22 **climate.** The average condition of the weather at a place over a period of years as exhibited
23 by temperature, wind velocity, precipitation and other factors.

24
25 **climax vegetation.** The vegetation that would exist if it was undisturbed.

26
27 **Code of Federal Regulations.** A codification of the general and permanent rules published
28 in the Federal Register by the departments and agencies of the Federal government. The
29 Code is divided into 50 titles that represent broad areas subject to Federal regulation. It
30 is issued quarterly and revised annually.

31
32 **collapse block.** A gravity-induced collapsed rock unit resulting from the formation of sink
33 holes.

34
35 **colloid.** Refers to a suspended particulate in a colloidal suspension or to a particle that is
36 small enough that it could be placed in colloidal suspension. A colloidal suspension is a
37 dispersion of extremely fine particles in a fluid with the characteristic size of the
38 particles (in at least one dimension) being so small that the behavior differs from that of
39 more common suspensions. The maximum size range for particles which may be
40 regarded as "colloidal" is subject to some variation, depending on context and
41 application.

42
43 **complementary cumulative distribution function.** Mathematically, a complementary
44 cumulative distribution function is equal to one minus a cumulative distribution



1 function. A cumulative distribution function is the sum (or integral) of the probability of
2 those values or variables that are less than or equal to a specified value. Complementary
3 cumulative distribution functions are a graphical display of the probability (the ordinate)
4 that the value of the variable will be greater than a specified value (the abscissa). For
5 the WIPP, the complementary cumulative distribution function displays the probability
6 that 10,000-year cumulative radionuclide releases from the disposal system for the
7 scenarios considered will exceed calculated values. Radionuclide releases are
8 normalized as stipulated in 40 CFR Part 191, Appendix A, and the complementary
9 cumulative distribution function is compared to the quantitative release limits specified
10 in 40 CFR § 191.13(a).”

11
12 **complexing agents.** Complexing agents increase the solubility of the complexed substance;
13 the aqueous solubility. They are often referred to as complexants.

14
15 **complexants.** Complexing agents increase the solubility of the complexed substance; the
16 aqueous solubility. They are often referred to as complexing agents.

17
18 **compliance assessment.** The analysis conducted to determine compliance with the Individual
19 Protection requirements in 40 CFR § 191.15 and the Groundwater Protection
20 Requirements in Subpart C.

21
22 **computational model.** The computational model is the implementation of the mathematical
23 model. (See definition of mathematical model below). The implementation may be
24 through analytical or numerical means. Often the analytical solution is numerically
25 evaluated (for example, numerical integration or evaluation of complex functions);
26 hence, both solution techniques are typically coded on the computer. Consequently, the
27 computational model is often called a computer model.

28
29 **computer model.** A computer code or set of codes that are used together to make
30 calculations for a corresponding mathematical description for a process or system (for
31 example, movement of groundwater through a geologic formation).

32
33 **conceptual model.** A set of qualitative assumptions used to describe a system or subsystem
34 for a given purpose. At a minimum, these assumptions concern the geometry and
35 dimensionality of the system, initial and boundary conditions, time dependence, and the
36 nature of the relevant physical and chemical processes. The assumptions should be
37 consistent with one another and with existing information within the context of the
38 given purpose.

39
40 **conductivity.** A material property describing the ease with which something can be
41 transferred through that material. Examples of conductivity are electrical conductivity
42 (electricity), thermal conductivity (heat transfer), or hydraulic conductivity (groundwater
43 flow).
44



1 **confirmation.** For the purposes of this document, a term used to indicate support or
2 establishment of certainty and/or validity of models used in reference to specific
3 performance issues of the repository over any specific time frame of interest. In general,
4 laboratory and field experiments at the WIPP and elsewhere are conducted to provide
5 data in support of this type of activity, such as for the gas generation model.
6 Confirmation is used in a mechanistic sense and is not intended to have specific legal
7 implications (See Validation).

8
9 **consequence analysis.** The analysis used to examine the effects of processes and events on
10 performance of the disposal system. (See Performance Assessment).

11
12 **conservative assumption.** Assumptions that result in the overestimation rather than
13 underestimation, of any phenomenon that could contribute to the release of
14 radionuclides from the disposal system.

15
16 **Consultation and Cooperation Agreement.** An agreement that affirms the intent of the
17 Secretary of Energy to consult and cooperate with the State of New Mexico with respect
18 to State public health and safety concerns. The term "Agreement" means the
19 July 1, 1981, Agreement for Consultation and Cooperation, as amended by the
20 November 30, 1984, "First Modification," the August 4, 1987, "Second Modification,"
21 and the March 22, 1988, modification to the Working Agreement.

22
23 **contact-handled transuranic waste.** Packaged transuranic waste that has a measured
24 external surface dose rate less than 200 millirems per hour at the surface of the
25 container.

26
27 **containment.** The retention of radioactivity within prescribed boundaries, such as within a
28 waste package. In this document, containment refers to retention within a system to
29 prevent releases in unacceptable quantities or concentrations during handling and
30 storage or to containment within the disposal system.

31
32 **contamination.** Introduction of radionuclides into the repository's immediate surroundings
33 such as into the brine via a breach of the containment system.

34
35 **controlled area.** The controlled area means (1) a surface location, to be identified by passive
36 institutional controls, that encompasses no more than 100 square kilometers and extends
37 horizontally no more than 5 kilometers in any direction from the outer boundary of the
38 original location of the radioactive wastes in a disposal system; and (2) the subsurface
39 underlying such a surface location. (40 CFR § 191.12). For WIPP, the controlled area
40 is defined by the area withdrawn by the Land Withdrawal Act.

41
42 **convection.** The circulatory motion that occurs in a fluid at a nonuniform temperature. The
43 transfer of heat by automatic circulation of fluids.



Title 40 CFR Part 191 Compliance Certification Application

1 **corrosivity.** The tendency of a metal to deteriorate by chemical decomposition.

2
3 **creep.** A very slow, usually continuous, time-dependent movement of soil or rock; refers to
4 the geologic phenomenon experienced as the gradual flow of salt under differential
5 stress.

6
7 **creep closure.** Closure of underground openings, especially openings in salt, by plastic flow
8 of the surrounding rock under lithostatic pressure.

9
10 **Cretaceous.** The final period of the Mesozoic era (after the Jurassic and before the Tertiary
11 period of the Cenozoic era), thought to have covered the span of time between 135 and
12 66 million years ago; also, the corresponding system of rocks. It is named after the Latin
13 word for chalk (creta) because of the English chalk beds of this age.

14
15 **crustal processes.** Any processes that occur within the earth's crust. Usually refers to
16 igneous metamorphic or tectonic activity.

17
18 **crystallographic.** This term deals with the system of forms among crystals, their structures,
19 and physical properties.

20
21 **CSH gel.** A cementitious material containing calcium-silicate-hydrate gel.

22
23 **cubic law.** The cubic law states that for a given change in gradient of the hydraulic head, flow
24 through a fracture is proportional to the cube of the fracture opening.

25
26 **cueta.** A hill or ridge with a gentle slope on one side and a steep slope on the other, specif.
27 an asymmetric ridge (as in the SW U.S.) with one face (dip slope) long and gentle and
28 conforming with the dip of the resistant bed or beds that form it, and the opposite face
29 (scarp slope) steep or even cliff-like and formed by the outcrop of the resistant rocks, the
30 formation of the ridge being controlled by the differential erosion of the gently inclined
31 strata.

32
33 **Culebra.** Culebra Dolomite Member, also referred to as Culebra, is the lower of two rock
34 units of dolomite within the Rustler Formation formed in the late Permian Period
35 between approximately 258 and 245 million years ago. It is a brown, finely crystalline,
36 argillaceous dolomite or dolomitic limestone with solution cavities. The Culebra is the
37 principal water-bearing unit of the Rustler Formation and has been identified as the
38 likely pathway for release of radionuclides to the subsurface accessible environment.

39
40 **cultural resource sites.** Human-associated ruins of archaeological significance.

41
42 **Cumulative Distribution Function.** A cumulative distribution function is the sum (or
43 integral) of the probability of those values of variable that are less than or equal to a
44 specific value.



1 **curie.** A quantitative measure of radioactivity equal to 3.7×10^{10} disintegrations per second.

2
3 **curie load.** The number of curies in a particular location or in a container such as a
4 radioactive waste drum. The curie load of a waste drum may be broken down into the
5 alpha, beta, and gamma curie loads to describe the amount of total radioactivity that is
6 contributed by disintegrations yielding alpha, beta, or gamma radiation.

7
8 **cuttings.** During rotary drilling, material contained in the cylindrical volume created by the
9 cutting action of the drill bit. In terms of releases from the WIPP through inadvertent
10 human intrusion, the maximum potential waste volume that could form cuttings is equal
11 to the cross-sectional area of the drill bit multiplied by the repository thickness.

12
13 **Darcy's Law.** A mathematical equation that can be used to compute the quantity of water
14 flowing through an aquifer. $Q = KA(h_A - h_B/L)$. Q is the discharge, A is the cross-
15 sectional area, $h_A - h_B$ is the difference in the height of the water (hydraulic head), L is
16 the flow length, and K is a coefficient, which is dependent upon the nature of the porous
17 medium.

18
19 **daughter product.** A nuclide that results from radioactive decay. Thus radium-226 decays
20 to radon-220, which in turn decays to polonium-216. The radon is the daughter of the
21 radium, and polonium is its daughter.

22
23 **Davis Mountains.** A mountain range in Texas south of the WIPP site making up the southern
24 most part of the Capitan Reef.

25
26 **decay, radioactive.** The decrease in the number of radioactive nuclei present in a radioactive
27 material due to their spontaneous transmutation. Also, the transmutation of a
28 radionuclide into another nuclide by the emission of a charged particle.

29
30 **decommissioning.** Actions taken upon closure of the repository to reduce potential
31 environmental, health, and safety impacts, including repository sealing as well as
32 activities to stabilize, reduce, or remove radioactive materials or demolish surface
33 structures.

34
35 **decommissioning phase.** The term "decommissioning phase" means the period of time
36 beginning with the end of the disposal phase and ending when all shafts at the WIPP
37 repository have been sealed.

38
39 **decontamination.** With respect to operational activities, decontamination is the removal of
40 unwanted material (especially radioactive material) from the surface of or from within
41 another material.



1 **defense waste.** Nuclear waste deriving from the manufacture of nuclear weapons and the
2 operation of naval reactors. Associated activities such as the research carried on in the
3 weapons laboratories also produce defense waste.
4

5 **deformation.** Any change in the original form or volume of rock masses. Folding, faulting,
6 solid flow, and dissolution and subsequent collapse are examples of deformation.
7

8 **degradation.** The wearing down or away of a material or the earth's surface whether on a
9 large or small scale and the products it produces. Examples of degradation include the
10 wearing down of drill casing from corrosion, and the wearing down of organics in the
11 waste materials from microorganisms.
12

13 **Delaware Basin.** Those surface and subsurface features that lie inside the boundary formed
14 to the north, east and west of the disposal system by the innermost edge of the Capitan
15 Reef, and formed, to the South, by a straight line drawn from the southeastern point of
16 the Davis Mountains to the most southwestern point of the Glass Mountains.
17

18 **Delaware Mountain Group.** A set of three formations that underlie the Castile Formation at
19 the Los Medaños site. The uppermost of these is the Bell Canyon Formation.
20

21 **demography.** The statistical study of human population with response to sizes and density,
22 distribution, and vital statistics.
23

24 **density.** Mass per unit volume (that is, kilograms per cubic meter).
25

26 **denudation.** The sum total of processes that results in the general lowering of the land
27 surface.
28

29 **Design Basis Earthquake.** An earthquake that is the most severe design basis accident of
30 this type and that produces the vibratory ground motion for which safety class items are
31 designed to remain functional.
32

33 **Design Basis Tornado.** A tornado that is the most severe design basis accident of that type
34 applicable to the area under consideration.
35

36 **detritivores.** Organisms that eat organic debris.
37

38 **deviatoric stress.** Refers to a state of stress disequilibrium in the subsurface, in which at least
39 one of the three perpendicular (normal) axes used to characterize stress at a point differs
40 in magnitude from the other two. The stress tensor used to characterize subsurface
41 stress can be written as the sum of two tensors. The first tensor represents a hydrostatic
42 state with all shear stresses equal to zero. The components of the second tensor are
43 shear stresses - these are deviatoric stresses and are independent of hydrostatic stress.
44



1 **Devonian.** A period of the Paleozoic Era (after the Silurian and before the Mississippian),
2 covering the span of time between 400 and 345 million years ago; also, the
3 corresponding system of rocks. It is named after Devonshire, England, where rocks of
4 this age were first studied.

5
6 **Devonian Woodford Shale.** This is a Devonian-age geological marker about 15,600 feet
7 deep at the WIPP that separates rocks of the Silurian era from rocks of the Mississippian
8 era.

9
10 **Dewey Lake Redbeds.** A formation of the Permian Period that overlies the Rustler
11 Formation and is composed of reddish brown marine mudstone and siltstones
12 interbedded with fine grained sandstone.

13
14 **diapirism.** The formation of an intrusion or diapir that penetrates through overlying layers of
15 more brittle rock and domes the overlying rock cover. Diapirs are commonly formed
16 from mobile igneous intrusions or from plastic salt bodies. The development of narrow
17 necks at depth often gives diapirs a balloon-like form.

18
19 **diagenetic.** Pertaining to or caused by diagenesis; for example, a “diagenetic change”
20 resulting from compaction, a “diagenetic structure” (such as a stylolite) formed after
21 deposition, a “diagenetic deposit” (such as dolomitized limestone or one consisting of
22 manganese nodules), or a “diagenetic environment” of rock consolidation.

23
24 **diffusion, molecular.** Transport of material (particularly the spread of a contaminant)
25 through the random motion of molecules in a fluid or solid. Diffusion results in the
26 slow transfer of material from regions of high concentration to regions of lower
27 concentration even when there is no net fluid flow.

28
29 **dike.** A tabular body of igneous rock that cuts across the structure of adjacent rocks or cuts
30 massive rocks. Most dikes result from the intrusion of magma. Some dikes occur in a
31 set of numerous parallel structures identified as dike swarms. Present in the vicinity of
32 the WIPP site are lamprophyre dikes. (See Lamprophyre)

33
34 **discharge point (or area).** In groundwater hydraulics, the point (or area) where water comes
35 out of an aquifer onto the surface.

36
37 **dispersion.** Signifies the tendency for a sharp change in fluid properties (for example,
38 temperature, contaminant concentration) to become broader and less abrupt as flow
39 progresses. For example, if a volume of contaminated water is injected into an aquifer,
40 the contact between the contaminated and uncontaminated water will initially be sharply
41 defined by a very steep gradient in contaminant concentration. As the water flows away
42 from the injection point, however, natural variations in the rock will mean that
43 contaminated water will move quickly through some large pores while nearby smaller
44 pores will retain uncontaminated water for a longer period. This, and other similar



Title 40 CFR Part 191 Compliance Certification Application

1 effects, will cause contaminated and uncontaminated water to intermingle, resulting in
2 dispersion of the contaminant "front."
3

4 **disposal.** The term "disposal" means permanent isolation of transuranic waste from the
5 accessible environment with no intent of recovery, whether or not such isolation permits
6 the recovery of such waste. Disposal of waste in a mined geologic repository occurs
7 when the waste has been emplaced and all the shafts to the repository are sealed.
8

9 **disposal phase.** The term "disposal phase" means the period of time during which
10 transuranic waste is disposed of at the WIPP, beginning with the initial emplacement of
11 transuranic waste underground for disposal and ending when the last container of
12 transuranic waste is emplaced underground for disposal and the shafts are sealed.
13

14 **disposal room.** An excavated cavity in the WIPP underground in which transuranic waste
15 will be emplaced during disposal operations.
16

17 **disposal system.** The disposal system is any combination of engineered and natural barriers
18 that isolate transuranic waste after disposal. For the purposes of the Waste Isolation
19 Pilot Plant, this will include the combination of the repository/shaft system and the
20 controlled area.
21

22 **disposal system performance.** Disposal system performance is measured in the Performance
23 Assessment in Section 6.5 of the Compliance Certification Application. It is compared
24 with the EPA requirements in the complementary cumulative distribution function.
25

26 **dissolution.** (1) The process whereby a space or cavity in or between rocks is formed by the
27 solution of part of the rock material. (2) The solution of material by a solvent (for
28 example, dissolution of waste by brine at the WIPP).
29

30 **disturbed rock zone.** That portion of the geologic barrier in which the physical and/or
31 chemical properties may have changed significantly as a result of underground
32 construction activities.
33

34 **Dockum Group.** The Triassic Dockum Group consists of three formations. In ascending
35 order, they are, (1) the Tecovas Formation, which consists of up to 300 feet of red shale,
36 siltstone, and fine-grained sandstone, (2) the Santa Rosa Sandstone, which is composed
37 of 100 to 650 feet of red, brown, and gray sandstone, and (3) the Chinle Formation
38 equivalent, which consists of up to 1,300 feet of red, maroon, and purple shales and
39 siltstones with lenses of fine-grained red-to-gray sandstone. The group is present only as
40 a thin wedge in Eddy County, thickening to the east in Lea County and in Texas.
41

42 **dolomite.** A sedimentary rock consisting mostly of the mineral dolomite: $\text{CaMg}(\text{CO}_3)_2$. It is
43 commonly found with limestone.
44



1 **dome (breccia pipe).** A type of hill found near the Los Medaños site; under at least a few of
2 these hills lies a roughly cylindrical volume of breccia (rock reconstituted of coarse rock
3 fragments).
4

5 **dome, salt.** A diapiric or piercement structure with a central, nearly circular salt plug,
6 generally 1 to 2 kilometers in diameter, that has risen through the enclosing sediments
7 from a deep mother bed of salt. In the continental U.S., salt domes are located primarily
8 in the Gulf Coast states.
9

10 **dose.** A general term indicating the amount of energy absorbed per unit mass from incident
11 radiation.
12

13 **dose conversion factor.** A numerical factor used in converting radionuclide uptake (curies)
14 in the body to the resultant radiation dose or dose commitment (rem or man-rem).
15

16 **dose equivalent.** The product of absorbed dose and appropriate factors to account for
17 differences in biological effectiveness due to the quality of radiation and its spatial
18 distribution in the body; the unit of dose equivalent is the rem (sievert in System of
19 International units). (40 CFR § 191.12)
20

21 **downdip.** To move in the direction at which a rock layer or any planar feature is inclined
22 (dip) from the horizontal.
23

24 **drift.** A horizontal passageway in a mine away from the main shaft.
25

26 **drilling fluids.** A suspension fluid used in drilling. The fluid is pumped through the inside of
27 the drill pipe and out through the jets of the drilling bit. It is used to lubricate the
28 drilling bit, seal off porous zones, and to counter-balance the pressure of oil, gas, or
29 water. The fluids consist of various substances in a finely divided state among which
30 bentonite and barite are common. Oil may also be used as a base instead of water.
31

32 **Driving Force Ratio.** A ratio established to assess the effects of density gradient when
33 examining hydraulic effects of fluid density variations in the Salado interbeds and the
34 Culebra. It is given by driving force ratio = $(\Delta\rho \nabla E)/(\rho_f \Delta H_f)$, where ΔH_f is the
35 gradient of freshwater head, $\Delta\rho$ is the difference between actual fluid density and
36 reference fluid density, ρ_f is the density of freshwater, and ∇E is the gradient of
37 elevation.
38

39 **E1, E2.** These are potential human intrusion scenarios used in computer modeling for
40 compliance purposes.
41

42 **effective dose.** The sum over specified tissues of the products of the dose equivalent received
43 following an exposure of, or an intake of radionuclides into, specified tissues of the
44 body, multiplied by appropriate weighting factors. This allows the various tissue-



Title 40 CFR Part 191 Compliance Certification Application

1 specific health risks to be summed into an overall health risk. The method used to
2 calculate effective dose is described in Appendix B of 40 CFR Part 191. (40 CFR
3 § 191.12)
4

5 **effluent.** Wastewater or airborne emissions discharged into the environment.
6

7 **Eh.** The oxidation potential is a measure of the oxidizing or reducing tendency of a solution.
8 It is written as a reduction reaction using the standard hydrogen reaction, $E_h \text{ (volts)} = E_h^0$
9 $+ RT/nF \ln \text{ (oxidized species/reduced species)}$.
10

11 **elasticity.** The ability of rocks to recover in size and shape (no long-term changes in rock
12 properties) resulting from minor deformation (for example, minor deflection of
13 anhydrite beds, deformation of rock due to passage of seismic wave, etc.). The property
14 by virtue of which a body resists and recovers from deformation produced by force.
15

16 **electrochemical.** Refers to chemical effects or reactions that produce, are mediated by, or
17 significantly influenced by electrical currents or accumulations of electrical charge.
18

19 **electrolytes.** A substance that will provide ionic conductivity when dissolved in water or
20 when in contact with water. The term is also used to designate solutions of such
21 substances.
22

23 **electromechanical.** Mechanical processes or effects that are driven by or produce electricity.
24

25 **electron acceptors.** Atoms or molecules that will readily accept an additional electron to
26 form an ionic or covalent bond. The availability of such chemical species is necessary
27 for some bacterial processes involving the decomposition of complex organic molecules.
28

29 **electrophoresis.** Migration of colloidal, or other particles in a liquid suspension which
30 responds to applications of an electric field.
31

32 **empirical.** Methods that rely on experience or observation alone without regard for system
33 and theory. They are capable of being verified or disproved by observation or
34 experiments.
35

36 **emplacement.** At the WIPP, the placing of radioactive wastes in the repository.
37

38 **engineered alternatives.** Potential modifications to the design or operation of the WIPP or to
39 waste forms that, if adopted, will provide increased assurance that the WIPP will
40 perform in compliance with environmental protection and safety requirements.
41

42 **engineered barriers.** Backfill, seals, and any other man-made barrier components of the
43 disposal system.
44



Title 40 CFR Part 191 Compliance Certification Application

1 **enthalpy.** The sum of the internal energy (heat) generated or absorbed when a reaction takes
2 place.

3
4 **eolian.** Pertaining to the wind. Also spelled aeolian.

5
6 **EPA unit.** The EPA unit is a measure of radioactivity that has been normalized by the
7 inventory of the disposal system in question and by the release limits specified by the
8 EPA in Appendix B, Table 1, of 40 CFR Part 191. Specifically, a quantity of
9 radioactivity of any radionuclide (expressed in curies) may be expressed in
10 dimensionless EPA units by dividing it by the product of the waste unit factor and the
11 EPA release limit for that radionuclide. The containment requirements of 40 CFR
12 § 191.13(a) establish release limits for the disposal system in terms of the probability
13 that 10,000 year cumulative releases will exceed 1 and 10 EPA units.

14
15 **epicenter.** The point on the earth's surface directly above the focus of an earthquake.

16
17 **equilibrium.** A state of balance between opposing forces or actions that is either static (as in
18 a body acted on by forces whose resultant is zero) or dynamic (as in a reversible
19 chemical reaction when the velocities in both directions are equal).

20
21 **erosion.** The processes whereby earthy or rock material is loosened or dissolved and removed
22 from any part of the earth's surface. It includes the processes of weathering, solution,
23 corrosion, and transportation.

24
25 **estuarine.** The seaward end of the widened tidal mouth of a river where fresh water comes
26 into contact with sea water.


27
28 **evaporite.** A sedimentary rock composed primarily of minerals produced by precipitation
29 from a solution that has become concentrated by the evaporation of a solvent, especially
30 salts deposited from a restricted or enclosed body of seawater or from the water of a salt
31 lake. In addition to halite (NaCl) these salts include potassium, calcium, and
32 magnesium chlorides and sulfates.

33
34 **event.** A phenomenon that occurs instantaneously or within a short time interval relative to
35 the time frame of interest.

36
37 **excavation.** The action of cutting, digging, or mining a cavity in the earth and the removal of
38 that mined material. The WIPP repository excavated in solid salt rock is an example of
39 excavation.

40
41 **exothermic.** The release of heat from a (chemical) reaction.

42
43 **extraction.** The withdrawal of natural resources with no provision for replenishment.



1 **fast Langrangian analysis of continua.** A two-dimensional finite-difference code used for
2 the WIPP to estimate subsidence above the repository for conditions ranging from no
3 backfill to emplacement of a highly compacted crushed salt backfill.
4

5 **fault.** A surface or zone of rock fracture along which there has been displacement.
6

7 **fault tree.** A tree-like cause-and-effect diagram of hypothetical events. Analysis of fault
8 trees is used to investigate failures in a system or concept.
9

10 **fauna.** An entire animal population of a given region, environment, formation, or time span.
11

12 **features, events, and processes.** Features, events, and processes that are potentially
13 important to long-term performance of the disposal system. A comprehensive set of
14 features, events, and processes relevant to the WIPP was considered in applying a
15 screening methodology to evaluate compliance with the numerical performance
16 requirements provided in 40 CFR Part 191.
17

18 **Federal Facilities Compliance Act.** An amendment, promulgated in 1992, to the Solid
19 Waste Disposal Act. Title I of the act grants the EPA administrative enforcement
20 authority against any department, agency, or instrumentality of the executive, legislative,
21 or judicial branch of the federal government. In regard to mixed wastes, sovereign
22 immunity for federal agencies is waived, consistent with a schedule provided in the act.
23 In addition, the act requires that the DOE prepare an inventory of mixed wastes and
24 mixed waste treatment capacities and technologies. For those mixed wastes for which
25 treatment capacities or technologies do not exist, the Department must prepare plans for
26 the development of the capacities or technologies.
27

28 **Fick's Law.** Mathematical equation in which steady state diffusion through solids is
29 represented.

$$J = -D \frac{\partial c}{\partial x}$$

30 J is the flux or diffusion current density, D is the diffusion coefficient, c is the volume
31 concentration of atoms, and x is the distance along the direction in which diffusion
32 occurs.
33

34 **filter bank.** An arrangement of air filters in series and/or parallel.
35

36 **final safety analysis report.** A safety document providing a concise but complete description
37 and safety evaluation of the site, the design, normal and emergency operations, potential
38 accidents, and predicted consequences of such accidents, and the means proposed to
39 prevent such accidents or to mitigate the consequences of such accidents. A final safety
40 analysis report documents the adequacy of safety analysis for a nuclear facility to ensure



2 that the facility can be constructed, operated, maintained, shut down, and
3 decommissioned safely and in compliance with applicable laws and regulations.

4 **fissile material.** Fissile material means any material consisting of or containing one or more
5 fissile radionuclides. Examples of fissile radionuclides are plutonium-238, plutonium-
6 239, plutonium-241, uranium-233, and uranium-235.

7
8 **fission.** The splitting of a heavy nucleus into two approximately equal parts, each the nucleus
9 of a lighter element, accompanied by the release of a large amount of energy and
10 generally one or more neutrons. Fission can occur spontaneously, but it usually follows
11 the absorption of neutrons.

12
13 **fissionable.** A nuclide that is available to undergo fission on absorption of a neutron with
14 energy over some threshold energy.

15
16 **flowpath.** The path traveled by a zero-charged, floating particle released into a groundwater
17 flow field.

18
19 **fluid.** A substance that is able to flow freely as liquids and gases do.

20
21 **fluvial.** Produced by or pertaining to a river or rivers. Applies to erosion (fluvial erosion) by
22 rivers or streams and deposition (fluvial deposition) by the accumulation of sediments in
23 a depositional environment associated with rivers or streams.


24
25 **flux.** The rate of transfer of heat, mass, electricity, groundwater, etc., across a unit surface
26 area in a unit time.

27
28 **formation (geologic).** The basic rock-stratigraphic unit in the local classification of rocks. It
29 consists of a body of rock (usually sedimentary) generally characterized by some degree
30 of internal lithologic homogeneity or distinctive features.

31
32 **Forty-niner Member.** Upper most member of the Rustler Formation formed in the Late
33 Permian Period, between approximately 250 and 245 million years ago. It consists of
34 upper and lower sulfates separated by claystone or halite. The member is about 60 feet
35 (18 meters) thick around the WIPP site.

36
37 **fracture porosity.** Porosity resulting from the presence of openings or fractures produced by
38 the breaking or shattering of an otherwise less porous rock.

39
40 **fractures.** A general term for any break in a rock, whether or not it has an associated
41 displacement. Fractures are the result of mechanical failure by stress. Fractures include
42 cracks, joints, and faults. Fracture porosity is the resulting opening produced by the
43 breaking or shattering of an otherwise less pervious rock.
44



1 **fugacities.** The vapor pressures of vapors assuming them to behave as ideal gases.

2
3 **galvanic coupling.** Refers to the establishment of an electrical current through chemical
4 processes. Galvanic coupling could lead to the establishment of potential gradients
5 between metals in the waste form, canisters, and other metals external to the waste form.
6

7 **gamma rays.** (X-rays) Short wavelength electromagnetic radiation emitted in the radioactive
8 decay of certain nuclides. Gamma rays are the same as gammas or gamma particles.
9

10 **gas generation.** Three gas generation processes are expected to be factors in the degradation
11 of transuranic waste in the WIPP repository. The generation of gaseous species is
12 expected to occur through chemical (that is, corrosion), microbial, and radiolytic
13 processes.
14

15 **gas generation model.** A computational model that can simulate and/or predict the rate and
16 quantity of gases generated by waste transformation processes in a disposal room of the
17 decommissioned repository.
18

19 **gas generation rate.** The combined gas production rate from all species of gases produced as
20 a result of transuranic waste transformations such as corrosion, and/or microbial
21 degradation, at any given time. The rate of gas production throughout the history of the
22 repository is expected to vary depending on repository conditions with respect to
23 humidity, total or partial brine inundation, competitive reactions that absorb specific
24 gases, and the ability of the repository to retain the gases generated. The term is also
25 applied to individual gases.
26

27 **gastropod.** Any of a large class (Gastropoda) of mollusks with a univalve shell or none
28 usually with a distinct head bearing sensory organs.
29

30 **Gatuña Formation.** A geologic formation covering the Permian Dewey Lake Formation, or
31 the Santa Rosa Formation of the late Pliocene Period. Considered to be fluvial in origin,
32 the Gatuña Formation varies in thickness and is overlain by the Mescalero Caliche.
33

34 **generator and/or storage sites.** Refers to the DOE sites nationwide where transuranic
35 wastes are generated and/or stored as a result of activities associated with nuclear
36 weapons production.
37

38 **geochemistry.** The study of the relative and absolute abundances of elements found on the
39 earth, and the chemical distribution and migration of individual elements located in
40 various parts of the earth (the atmosphere, hydrosphere, etc.).
41

42 **geomechanical.** The response of natural earth materials to the application of deforming
43 forces. These forces would include the excavation and subsequent deformational
44 response of the WIPP disposal rooms and drifts.



1 **geomorphology.** The description and interpretation of landforms on the earth's surface.

2
3 **geophysical log.** A graphic record of the measured or computed physical characteristics of
4 the rock section encountered in a well, plotted as a continuous function of depth.
5 Measurements are made by a sonde as it is withdrawn from the borehole by a wireline.
6 Several measurements are usually made simultaneously, and the resulting curves are
7 displayed side by side on the common depth scale. Both the full display and the
8 individual curves are called logs. Well logs are commonly referred to by generic type.

9
10 **geophysical techniques.** The methods used to obtain geophysical data (that is, values of the
11 physical parameters of the earth). The methods of techniques generally involve the
12 study of ambient fields (gravitational, magnetic, electrical, temperature) or the study of
13 the effects of applied fields (seismics through the application of shock waves, electrical
14 through the direct application of electrical currents — resistivity methods — and by
15 electromagnetic induction.)

16
17 **geostatistics.** A branch of statistics used to describe and interpret spatial data.

18
19 **geothermal.** The internal energy of the earth, available from heat sources produced at depth
20 within the earth's crust.

21
22 **getters.** Substances that sorb gases, such as carbon dioxide (CO₂), and may be added with
23 other materials to mitigate the pressure buildup in the repository and radionuclide
24 mobility.

25
26 **Gibbs Free Energy.** A thermodynamic quantity defined as $H - TS$. H is enthalpy, S is
27 absolute entropy, and T is temperature. The magnitude of the change in Gibbs Free
28 Energy for a reaction determines whether it is reversible or will occur spontaneously at
29 given temperature and pressure.

30
31 **glaciation.** Applies to both the formation, progression, and recession of glaciers and the
32 resulting alteration of the earth's ground surface through erosion and deposition.

33
34 **glove box.** A sealed box in which workers, remaining outside and using gloves attached to
35 and passing through openings in the box, can safely handle and work with radioactive
36 materials under controlled conditions.

37
38 **Goat Seep.** The Goat Seep was a reef, which grew primarily upward and formed a barrier
39 around a considerable portion of the western side of the Delaware Basin. The lower
40 portion of the Goat Seep is thick-bedded, and the upper portion is a massive light gray,
41 fine-crystalline to saccharoidal, and in places very porous, dolomite.

42



1 **graben.** An elongate, relatively depressed crustal unit or block that is bounded by roughly
2 parallel faults on its long sides. It is a structural form that may or may not be
3 geomorphologically expressed as a rift valley.
4

5 **gradient.** Spatial changes in the value of a variable. Changes are along vertical, inclined or
6 horizontal distances, such as the groundwater pressure gradient of the Culebra Dolomite
7 changing in the vicinity of the WIPP site.
8

9 **greenhouse effect.** Warming of the earth's surface and the lower layers of atmosphere that
10 tends to increase with increasing atmospheric carbon dioxide and other gases, and is
11 caused by conversion of solar radiation into heat.
12

13 **groundwater.** Water below the land surface in a zone of saturation. (40 CFR § 191.12)
14

15 **grout.** A mortar of cement slurry (of high water content) used to plug potential fluid-flow
16 paths in geologic or engineered structures.
17

18 **Guadalupean.** Geological group of rocks below the Castile about 4,100 feet to about 8,000
19 feet below the surface at the WIPP. Contains the Bell Canyon, Brushy Canyon, and
20 Cherry Canyon formations.
21

22 **gypsite.** An earthy variety of gypsum containing dirt and sand, found only in arid regions.
23

24 **gypsum.** A mineral consisting of hydrous calcium sulfate: $\text{CaSO}_4 \cdot 2\text{H}_2\text{O}$. It is soft and,
25 when pure, white.
26

27 **half-life.** The time required for the activity of a group of identical radioactive nuclei to decay
28 to half its initial value. (See definition for biological half-life).
29

30 **halite.** The mineral rock salt: NaCl .
31

32 **halophilic.** A type of organism that flourishes in a salty environment such as salt lakes and
33 marshes. Halophilic microorganisms may be present in WIPP's disposal environment at
34 the time of closure.
35

36 **halotolerant.** A type of organism that is tolerant of salty environments but does not prefer
37 them. Halotolerant microorganisms may be introduced into the waste before disposal
38 from semi-salty environments such as the skin of humans.
39

40 **health physics.** The science concerned with the recognition, evaluation, and control of health
41 hazards from ionizing radiation.
42

43 **heavy metal.** All uranium, plutonium, or thorium placed into a nuclear reactor. (40 CFR
44 § 191.12)



1 **HEPA filter.** A high-efficiency particulate air filter usually capable of 99.7 percent efficiency
2 as measured by a standard photometric test using 0.3-micron droplets (aerodynamic
3 equivalent diameter) or dioctylphthalate.
4

5 **high-level waste.** Radioactive waste resulting from the reprocessing of spent fuel. Discarded,
6 unprocessed spent fuel is also high-level waste. It is characterized by intense,
7 penetrating radiation and by high heat-generation rates. Even in protective canisters,
8 high-level waste must be handled remotely.
9

10 **horizon.** In geology, an interface indicative of a particular position in a stratigraphic
11 sequence. For instance, the waste-emplacement horizon in the Salado Formation at the
12 WIPP is the level about 650 meters (2,150 feet) deep where openings are mined for
13 waste disposal.
14

15 **host rock.** The rock unit, in this case the Salado Formation, in which the radioactive waste is
16 to be emplaced.
17

18 **hot cell.** A heavily shielded compartment in which highly radioactive material can be
19 handled, generally by remote control.
20

21 **human intrusion.** (See Inadvertent Human Intrusion).
22

23 **humics.** Matter that is derived from humus, which is a brown or black complex of variable
24 material resulting from partial decomposition of plant or animal matter and forming the
25 organic portion of soil.
26

27 **hummocky.** Containing rounded knolls or small hills.
28

29 **hydration.** The uptake of water by a substance, generally involving reactions in which the H-
30 OH bond of the water molecules is not broken. (See hydrolysis.)
31

32 **hydraulic conductivity.** A quantity defined in the study of groundwater hydraulics that
33 describes the ability of rock to transmit groundwater. It is measured in feet per day or
34 equivalent units. It is equal to the hydraulic transmissivity divided by the thickness of
35 the aquifer.
36

37 **hydraulic gradient.** A quantity defined in the study of groundwater hydraulics that describes
38 the rate of change of hydraulic head with distance.
39

40 **hydraulic potential (or hydraulic head).** Hydraulic pressure corrected for the potential
41 energy of elevation. In an aquifer it is equivalent to the highest level of a column of
42 water that the pressure in the aquifer will support. It is measured relative to a specified
43 level, which in this document is the mean sea level.
44

1 **hydraulic transmissivity.** A rate at which water is transmitted through a unit width of
2 aquifer. It is measured in square meters per second or equivalent units. (See
3 transmissivity).
4

5 **hydraulic transport.** The transport of dissolved or suspended substances by groundwater.
6

7 **hydraulics, hydrology.** These two terms tend to be used interchangeably, but they are not the
8 same. Hydraulics is an engineering discipline; hydrology is the related science.
9 Hydraulics deals with the flow of water. Hydrology deals with water: its properties,
10 circulation, and distribution, from the time it falls as rainwater until it is returned to the
11 atmosphere through evapotranspiration or flows into the ocean.
12

13 **hydrocarbon.** An organic compound (as acetylene or benzene) containing only carbon and
14 hydrogen and often occurring in petroleum, natural gas, coal, and bitumens.
15

16 **hydrolysis.** A chemical reaction between water and another material in which two or more
17 new species are formed, implying that the H-OH bond in the water molecules is broken.
18

19 **hydrostatic.** The pressure and equilibrium of liquids such as the pressure exerted by water at
20 any given point in a body of water at rest. The hydrostatic pressure of groundwater is
21 generally due to the weight of water in higher levels in the zone of saturation.
22

23 **igneous.** Rocks that are formed by solidification from a molten or partially molten state. For
24 convenience, igneous rocks are divided into two major classes, plutonic and volcanic,
25 which describe igneous rocks formed at depth and at or near the land surface,
26 respectively.
27

28 **inadvertent human intrusion.** The accidental violation of the disposal system through
29 human activity such as mining or exploration drilling. Inadvertent and intermittent
30 intrusion by drilling for resources (other than those resources provided by the waste in
31 the disposal system or engineered barriers designed to isolate such waste) is the most
32 severe human intrusion scenario (40 CFR § 194.33[b][1]).
33

34 **infiltration.** The penetration of water through the ground surface into subsurface soil.
35

36 **ingestion.** The uptake of materials taken into the body by way of the digestive tract.
37

38 **ingrowth.** The process in which a nuclide results from the radioactive disintegration of a
39 radionuclide. The nuclide can be formed directly or as the results of successive
40 transformations in a radioactive series. A decay product may be either radioactive or
41 stable.
42

43 **injection well.** A well into which fluids are injected.
44





1 **in situ.** In the natural or original position. The phrase is used in this document to distinguish
2 in-place experiments, rock properties, and so on, from those measured in the laboratory.

3
4 **insoluble.** Unable to be dissolved.

5
6 **intensity, earthquake.** A measure of the effects of an earthquake on humans and structures
7 at a particular place. Not to be confused with magnitude.

8
9 **interbed.** A usually thin layer of rock, typically of one kind of rock material, occurring
10 between or alternating with layers of another rock type. The Salado Formation in the
11 vicinity of the WIPP site contains layers of anhydrite that are interbedded between more
12 massive layers of halite.

13
14 **interfacial energy.** Energy associated with an interface between two materials (surfaces).
15 Changes in interfacial energy control effects such as the wetting of a solid surface by a
16 liquid.

17
18 **intergranular.** The porosity or spaces occurring between the grains of granular or crystalline
19 rocks in which the void space usually contains a liquid.

20
21 **interpreted data sets.** Data sets derived from a surrogate data source where direct
22 measurement is not possible.

23
24 **interstitial brine.** Brine distributed in the pore space (voids) of a rock mass.

25
26 **intragranular.** The porosity or spaces occurring within a grain or crystal, especially within
27 skeletal material of a carbonate sedimentary rock in which the void space may or may
28 not contain a liquid.

29
30 **intraplate rifting.** Rifting of the continental or oceanic crust within a crustal plate Typically
31 applies to areas that have shown regional extensional tectonic activity away from the
32 plate boundaries and where igneous rocks have formed intrusive sill, dikes, volcanic
33 plugs, and other intrusive bodies. Along the Rio Grande valley, 120 miles (200
34 kilometers) west of the WIPP site, intraplate rifting has taken place during the Tertiary
35 and Quaternary Periods. (See Rio Grande Rift).

36
37 **ion exchange.** A phenomenon in which chemical species in one phase or material exchange
38 with similar species in another phase.

39
40 **isopach.** A line drawn on a map through points of equal true thickness of a designated
41 stratigraphic unit or group of stratigraphic units.

42
43 **isotope.** Any of two or more species of atoms of a chemical element with the same atomic
44 number but with differing atomic mass.



Title 40 CFR Part 191 Compliance Certification Application

1 **Jurassic.** The second period of the Mesozoic era (after the Triassic and before the
2 Cretaceous), thought to have covered the span of time between 190 and 135 million
3 years ago; also, the corresponding system of rocks.
4

5 **K-Ar.** Determination of the age of a mineral or rock in years, based on measurement of the
6 ratio of radiogenic argon-40 to potassium-40 and the known radioactive decay rate of
7 potassium-40 to argon-40.
8

9 **karst.** A type of topography that is formed on limestone, gypsum, and other rocks by
10 dissolution, and that is characterized by sinkholes, caves, and underground drainage.
11

12 **K_d.** A symbol used to designate a coefficient used in calculating chemical retardation.
13

14 **kelvin.** The name of the absolute temperature scale generally used in scientific measurements
15 and calculations. The unit of temperature is also designated a kelvin.
16

17 **kinetic.** An adjective referring to motion and, by extension, to changing conditions. In
18 chemistry, kinetic behavior refers to the performance of a system in which ongoing
19 chemical reactions are changing the quantities or character of the substances present in
20 the system at a certain rate and are thereby moving towards equilibrium.
21


22 **lacustrine.** The term is used to identify sediments formed by deposition in lakes and to
23 identify chemical and biological processes that typically occur in lakes.
24

25 **Lamar Limestone.** A limestone member at the top of the Bell Canyon Formation, known as
26 the Lamar limestone, is recognizable over a considerable part of the Delaware Basin.
27


28 **lamprophyre.** A group of dark-colored intermediate igneous rocks characterized by
29 unrestricted crystal structures and a high percentage of mafic minerals (that is, biotite,
30 hornblende, and pyroxene). Lamprophyre igneous rocks commonly occur in the form of
31 dikes. (See Dike) This type of dike is present in the vicinity of the WIPP site and is
32 identified by the dark minerals that occur both as phenocrysts and in the groundmass and
33 light minerals occurring only in the groundmass.
34

35 **Land Withdrawal Act.** Public Law 102-579, which withdraws the land at the WIPP site
36 from "entry, appropriation, and disposal"; transfers jurisdiction of the land from the
37 Secretary of the Interior to the Secretary of Energy; reserves the land for activities
38 associated with the development and operation of the WIPP; and includes many other
39 requirements and provisions pertaining to the protection of public health and the
40 environment.
41

42 **langbeinite.** A mineral, $K_2Mg_2(SO_4)_3$, used in the fertilizer industry as a source of potassium
43 sulfate present in economic quantities in the McNutt Potash Zone.
44



- 1 **Latin hypercube sampling.** A Monte Carlo sampling technique that divides the range of
2 each variable into intervals of equal probability and samples from each interval.
3
- 4 **leachate.** Means any liquid, including any suspended components in the liquid, that has
5 percolated through or drained from waste.
6
- 7 **leaching.** The process of extracting a soluble component from a solid by the percolation of a
8 solvent (in this report, water) through the solid.
9
- 10 **Leonardian.** The geologic formation from 8,000 feet to 11,400 feet below the surface at the
11 WIPP. Middle of the Permian Period.
12
- 13 **level-line survey.** A cross-country survey in which changes in elevation with respect to sea
14 level are very carefully measured.
15
- 16 **ligands.** A molecule, ion, or atom that is attached to the central atom of a coordination
17 compound. Ligands are also called complexing agents.
18
- 19 **lithofacies.** A lateral, mappable subdivision of a designated stratigraphic unit, distinguished
20 from adjacent subdivisions on the basis of lithology, including all mineralogic and
21 petrographic characters and those paleontologic characters that influence the appearance,
22 composition, or texture of the rock.
23
- 24 **lithology.** The study and examination of the physical characteristics (color, mineralogic
25 composition, grain size, etc.) of rocks, generally determined from a hand specimen with
26 the aid of a low-powered magnifier.
27
- 28 **lithosphere.** The outer solid shell of the earth considered to be about 50 miles (80 kilometers)
29 in thickness. Composed of the rocks that include both the earth's crust and upper
30 mantle.
31
- 32 **lithostatic pressure.** The vertical pressure at a point in the Earth's crust, equal to the pressure
33 caused by the weight of the overlying rocks.
34
- 35 **Livingston Ridge.** Topographic feature marking the eastern boundary of Nash Draw.
36
- 37 **long term.** Refers to the 10,000 years after shaft sealing for which performance assessment
38 calculations and models assess the behavior of the repository with respect to compliance
39 with 40 CFR Part 191 and 40 CFR § 268.6.
40
- 41 **Los Medaños.** In this report, the area in southeastern New Mexico surrounding the site
42 proposed for the WIPP repository. In Spanish it means "dune country."
43



1 **lower explosive limit.** The minimum concentration of gas or vapor in air below which a
2 substance does not burn when exposed to an ignition source.

3
4 **macromolecule.** A very large molecule (such as proteins, cellulose, and rubbers) that
5 contains hundreds or thousands of atoms.

6
7 **Magenta Dolomite Member.** The upper of two layers of dolomite within the Rustler
8 Formation that are locally water-bearing. Also called Magenta Member.

9
10 **magma.** Naturally occurring molten rock material which is generated within the earth. This
11 material is capable of intrusion and extrusion, from which igneous rocks are derived
12 through solidification and related processes.

13
14 **magnitude, earthquake.** A measure of the total energy released by an earthquake. Not to be
15 confused with intensity.

16
17 **Malaga Bend.** A sharp bend in the Pecos River 20 miles southeast of Carlsbad, New Mexico,
18 and directly east of the town of Malaga.

19
20 **man-rem.** A unit of population dose.

21
22 **marker beds.** Marker beds are well defined layers of rock that mark distinct divisions in
23 major geological strata or geological time frames.

24
25 **mathematical model.** In the context of the WIPP, mathematical formulations developed to
26 represent the processes at the WIPP site. The conceptual models provide the context
27 within which these mathematical models must operate and define the processes they
28 must characterize. The mathematical models are predictive in the sense that, once
29 provided with the known or assumed properties of the system and possible perturbations
30 to the system, they predict the response of the system. The processes represented by
31 these mathematical models include fluid flow, mechanical deformation, radionuclide
32 transport in groundwater, and removal of waste through intruding boreholes.

33
34 **matrix porosity.** The porosity of the matrix or finer parts of a rock, as opposed to the
35 porosity imparted to a rock through fractures, vugs, and voids.

36
37 **maximally exposed person.** A hypothetical person who is exposed to a release of
38 radioactivity in such a way that the person receives the maximum possible individual
39 dose or dose commitment. For instance, if the release is a puff of contaminated air, the
40 maximally exposed person is the individual at the point of largest ground-level
41 concentration who stays there during the whole time of the cloud passage. The use of
42 this term is not meant to imply that there is such a person, but only that thought is being
43 given to the maximum exposure a person could receive.



1 **maximum individual dose.** The highest dose delivered to the whole body or to an individual
2 organ that a person can receive from a release of radioactivity. The hypothetical person
3 who receives this dose, the maximally exposed person, is one whose location and
4 activities maximize the dose. For instance, the person may be at the point of maximum
5 concentration of a radioactive cloud for the whole time it takes to pass.
6

7 **MB139.** A marker bed located 1 meter below the WIPP site and is on average one meter
8 thick. It consists mostly of anhydrite and clay.
9

10 **mCi.** Abbreviation for millicurie. One-thousandth (0.001) of a curie.
11

12 **McNutt Potash Zone or Member.** The middle member of the Salado Formation measuring
13 348 to 413 feet (106 to 126 meters) thick in the vicinity of the WIPP site. This member
14 contains reddish-orange and brown halite with deposits of sylvite and langbeinite from
15 which potassium salts are mined. The McNutt Potash Member was deposited in the
16 Late Permian Period between approximately 258 and 250 million years ago.
17

18 **mean.** The average value. For a given set of n values, the mean is the sum of their values
19 divided by n .
20

21 **median.** The median of a set of data is the value such that half of the observations are less
22 than that value and half are greater than that value.
23

24 **megapascal.** 10^6 pascals.
25

26 **Mercalli intensity.** A scale of measurement of earthquake intensity.
27

28 **Mescalero Caliche.** An informal name for a layer of caliche of varying thickness found in the
29 WIPP area.
30

31 **Mescalero Plain.** The Plain is composed of the ground surface east of the Pecos River valley
32 and is one of the principal geomorphic features associated with the WIPP site. It
33 consists of a poorly drained surface covered by gravels, eolian sand, and caliche. The
34 plain has developed since the Early to Middle Pleistocene time (approximately 1.6
35 million year ago).
36

37 **Mesozoic.** An era of geologic time from the end of the Paleozoic Era, 245 million years ago,
38 to the beginning of the Cenozoic era, 66 million years ago. It comprises the Triassic,
39 Jurassic, and Cretaceous period.
40

41 **metabolized.** Refers to a material which has been consumed or transformed by the life
42 processes of an organism.
43



1 **metamorphic.** Metamorphism is changes in rocks brought on by the effects of temperature,
2 pressure and/or the introduction of new chemical substances. It requires deep burial
3 below the zones of weathering and cementation.
4

5 **methanogenesis.** The generation of methane through the decomposition of organic matter in
6 wastes.
7

8 **microbial.** The action of microbes or microorganisms. In the context of the WIPP, the term
9 is mainly applied to the waste degradation and gas generation activity of
10 microorganisms.
11

12 **mineralization.** The process whereby minerals are precipitated out of a solution into a porous
13 or fracture zone. For example, if the Salado Formation brine is transported to the
14 Culebra Dolomite cooling may cause the precipitation of salt minerals.
15

16 **Miocene.** The fourth of five epochs into which the Tertiary Period is divided; also, the
17 corresponding system of rocks. It is subsequent to the Oligocene Epoch and precedes the
18 Pliocene Epoch (approximately 23.7 to 5.3 million years ago).
19

20 **miscellaneous hazardous waste management unit.** A waste management unit where
21 hazardous waste is treated, stored, or disposed of, and that is not a container, tank,
22 surface impoundment, pile, land treatment unit, landfill, incinerator, boiler, industrial
23 furnace, underground injection well, or unit eligible for a research, development, and
24 demonstration permit (40 CFR § 260.10).
25

26 **Mississippian.** Third youngest period of the Paleozoic Era and the corresponding system of
27 rocks. Prior to the Pennsylvanian and representing approximately the interval from 360
28 to 320 million years ago. Mississippian rocks are present from 15,000 to 15,600 feet
29 below the surface at the WIPP.
30

31 **Mississippian Limestone.** The limestone is light gray to brown, finely crystalline and
32 commonly cherty, with a basal dark gray organic-rich shale unit. The limestone partially
33 grades to shale southeastward from the northern margin of the Delaware Basin.
34

35 **mixed waste.** Mixed waste contains both radioactive and hazardous components, as defined
36 by the Atomic Energy Act and the Resource Conservation and Recovery Act,
37 respectively.
38

39 **model.** An investigative technique using a mathematical or physical representation of a
40 system that accounts for all or some of its known parameters. Models are often used to
41 test the effect of changes of system components on the overall performance of the
42 system.
43



1 **model validation.** The process of ensuring (through sufficient testing of a model using actual
2 site data), that a conceptual model, and corresponding mathematical and computer
3 models, correctly simulate a physical process with sufficient accuracy.
4

5 **model verification.** The process of ensuring (for example, through tests on ideal problems)
6 that a computer code (computational model) correctly performs the necessary functional
7 operations (such as solving the mathematical model). Given that a computer code
8 correctly solves the mathematical model, the physical assumptions of the mathematical
9 model must then be checked through validation.
10

11 **molality.** The molality of a solution is the number of moles of solute in 1,000 g of solvent. A
12 solution which contains one mole of solute in 1,000 g of solvent is called a one-molal
13 solution.
14

15 **molarity.** The molarity of a solvent is the number of moles of solute per liter of solution. A
16 solution containing one gram formula weight, or mole, of the solute in one liter of
17 solution is a one-molar solution.
18

19 **Monte Carlo Analysis/Technique.** A technique that obtains distribution of approximate
20 solutions to a problem by using statistical sampling techniques and computer
21 simulations. For the WIPP performance assessment, the method is used to evaluate the
22 distribution of the consequence results and thereby approximate the uncertainty in the
23 results.
24

25 **Morrow Formation.** This formation was formed in the Early Pennsylvanian Period
26 approximately 320 million years ago. It consists of fine to coarse grain sandstones with
27 dark gray shales and some thin limestones layers present in the upper section. It is
28 approximately 14,000 feet (4,200 meters) below the surface at the WIPP and 1,200 feet
29 (366 meters) in thickness. The Morrowan sandstones are a source of natural gas
30 production north of the WIPP site.
31

32 **MTHM.** Metric tons of heavy metal; used in the release limit described in Appendix A of 40
33 CFR Part 191.
34

35 **mudstone.** An indurated mud having the texture and composition of shale, but lacking its
36 fine lamination or fissility; a blocky or massive, fine-grained sedimentary rock in which
37 the proportions of clay and silt are approximately equal; a nonfissile mud shale.
38

39 **multi-mechanism deformation model.** A simulation model developed to help predict the
40 behavior, particularly the rate of room closure, of WIPP underground openings.
41

42 **Nash Draw.** A shallow valley, approximately five miles wide, open to the southwest located
43 to the west of the WIPP site.
44



1 **natural background radiation.** Radiation in the human environment from naturally
2 occurring elements and from cosmic radiation. See background (radiation).

3
4 **natural barriers.** The repository host rock and surrounding geologic structures and
5 formations. The natural barriers extend from the engineered barrier to the compliance
6 boundary.

7
8 **near future.** Defined for any activity that is initiated prior to shaft sealing (preclosure), even
9 if the completion will occur after shaft sealing (postclosure). The demarcation between
10 near future and future is the date of shaft sealing. All activities initiated after shaft
11 sealing are "future" activities. Any activity initiated prior to shaft sealing is a "near
12 future" activity and is part of the undisturbed performance.

13
14 **neutron.** An elementary particle that has approximately the same mass as the proton but
15 lacks electric charge, and is a constituent of all nuclei having mass number greater than
16 one.

17
18 **New Mexico Bureau of Mines and Mineral Resources.** The New Mexico Bureau of Mines
19 and Mineral Resources is a division of the New Mexico Institute of Mine Technology.

20
21 **nitrates.** The salts from nitric acid generating sodium nitrate and potassium nitrate. These
22 salts are used in the agricultural industry as fertilizers. When used as a fertilizer, nitrates
23 can be a cause for groundwater chemistry changes.

24
25 **nondestructive assay.** The remote determination of the radionuclide content inside a waste
26 package.

27
28 **non-Salado.** In this report, Non-Salado refers to the hydrological aspects of all geologic
29 formations above and below the Salado Formation. It pertains to the conceptual models
30 used for flow and transport of radionuclides within these formations.

31
32 **nuclear criticality.** A self-sustaining nuclear chain reaction from sufficient mass of a
33 fissionable material.

34
35 **Nuclear Energy Agency.** The Nuclear Energy Agency was established on February 1, 1958
36 under the name of OEEC European Nuclear Energy Agency. Nuclear Energy Agency
37 membership today consists of all the European member countries of Organisation for
38 Economic Cooperation and Development as well as Australia, Canada, Japan, the
39 Republic of Korea, Mexico and the United States. The primary objective of Nuclear
40 Energy Agency is to promote cooperation among the governments of its participating
41 countries in furthering the development of nuclear power as a safe, environmentally
42 acceptable and economic energy source.



1 **nucleation.** The process by which the formation of a crystal or some other particulate from
2 the liquid state, vapor, or solution is begun. Often a minute particle of a foreign
3 substance provides the initial nucleus for crystal growth.

4
5 **nuclide.** A species of atom characterized by the number of protons and neutrons in its
6 nucleus.

7
8 **nuclide inventory (radionuclide inventory).** A list of the kinds and amounts of
9 radionuclides in a container or a source. Amounts are usually expressed in activity
10 units: curies or curies per unit volume.

11
12 **nutrient.** Applies to any substance assimilated by living things that promotes growth. The
13 term generally applies to nitrogen and phosphorus in waste water, but is also applied to
14 other essential and trace elements.

15
16 **Ochoan.** The Late Permian Ochoan series consists primarily of evaporites that were
17 deposited during recurrent retreats of a shallow sea restricted by the Guadalupian reefs.
18 The lower three formations in the series, the Castile, Salado and Rustler, comprise what
19 is perhaps the thickest and most extensive evaporite rock sequence in North America.

20
21 **Ogallala Formation.** Deposition of the Ogallala began about 12 million years ago, in
22 Miocene time. The Ogallala represents the first preserved sedimentary record in the
23 vicinity of the Delaware Basin since Cretaceous deposition. The Ogallala formed a
24 thick mantle throughout the Permian Basin, producing the even surface of the High
25 Plains, called the Llano Estacado in western Texas and eastern New Mexico. Locally,
26 eolian activity played a part in deposition, and periodically, widespread soils formed.

27
28 **one hundred-year storm.** A storm that, on a statistical basis, is expected to recur only once
29 every hundred years.

30
31 **oolite.** A sedimentary rock, usually a limestone, made up chiefly of ooliths cemented
32 together. Syn, roestone, eggstone. An oolith is a spherical or subspherical rock particle
33 that has grown by accretion around a nucleus. The commonest type of oolith is
34 calcareous, although primary ooliths of iron minerals are also found.

35
36 **order of magnitude.** A factor of ten. When a measurement is made with a result such as $3 \times$
37 10^7 , the exponent of 10 (here 7) is the order of magnitude of that measurement. To say
38 that this result is known to "within an order of magnitude" is to say that the true value
39 lies between (in this example) 3×10^6 and 3×10^8 .

40
41 **Ordovician.** The second oldest period of the Paleozoic Era and also the corresponding
42 system of rocks. It is represented by the rock units between 16,900 feet and 18,200 feet
43 below the surface in the vicinity of the WIPP. The Ordovician Period is subsequent to



1 the Cambrian Period and followed by the Silurian Period, between approximately 505
2 and 438 million years ago.

3
4 **orthogonal reactions.** Chemical reactions that do not affect or interfere with each other.

5
6 **osmotic.** The passage of pure liquid from a region of higher solute concentration to a region
7 of lower solute concentration, usually through some intervening solid or liquid barrier
8 such as a membrane, which restricts mixing. Thus, if a volume of dilute salt water is
9 separated from a volume of concentrated brine by a semi-permeable membrane, water is
10 passed by osmosis through the membrane from the dilute solution to the high-salinity
11 brine.

12
13 **overpack.** A container put around another container. In the WIPP, overpacks would be used
14 on damaged or otherwise contaminated drums, boxes, and canisters that it would not be
15 practical to decontaminate.

16
17 **oxic.** Describes a process that occurs in the presence of oxygen (O_2), such as the corrosion of
18 steels, iron-base alloys, and other metal wastes, using O_2 and water (H_2O) to produce
19 iron oxide and hydrogen gas (H_2).

20
21 **oxidation.** The chemical process whereby electrons are removed from an atom or ion.
22 Originally referring to chemical reactions involving oxygen, the term is also used to
23 designate any chemical reaction in which the reactant gives up one or more electrons.
24 Thus, when hydrogen (H_2) and chlorine (Cl_2) react to form $2H+Cl^-$, the hydrogen is
25 "oxidized" by the chlorine. The oxidation state of an ion is expressed as a positive or
26 negative number representing the ionic or effective charge.

27
28 **oxides.** Minerals in which cationic atoms are bonded to oxygen atoms.

29
30 **packaging.** The assembly of components necessary to ensure compliance with packaging
31 requirements. It may consist of one or more receptacles, absorbent materials, spacing
32 structures, thermal insulation, radiation shielding, and devices for cooling or absorbing
33 mechanical shocks. The vehicle, tie-down system, and auxiliary equipment may be
34 designated as part of the packaging.

35
36 **Paleozoic.** Major geological era from 245 million years to 570 million years. Denotes a wide
37 range of geological strata from different geological periods (that is, Permian,
38 Pennsylvanian, Mississippian, etc).

39
40 **panel.** A group of several underground rooms connected by drifts. Within the WIPP, a panel
41 consists of seven rooms connected by drifts at each end.

42
43 **PANEL.** Sandia National Laboratory computer code name for a program which simulates the
44 process of waste mobilization.



1 **parameter.** The underlying elements ($x = x_1, \dots, x_n, \dots, x_{nv}$) of a computational model. As x
2 changes so does the model result. The individual parameters, x_n , may be vectors,
3 tensors, higher order quantities, or even functions, but are usually scalar quantities.
4 Furthermore, the individual parameters are usually coefficients of the mathematical
5 model, but they may also relate to scenario uncertainty or model form uncertainty.

6
7 **Pascal.** A unit of pressure obtained by dividing force (in Newtons) by area (in meters
8 squared).

9
10 **passive/active neutron analysis.** An assaying technique used to determine the amount of
11 fissionable material in a container of waste. Background neutrons are counted as well as
12 neutrons from artificially-induced fissioning in the waste. The analysis assumes that the
13 isotopic composition is already known.

14
15 **passive institutional controls.** “(1) [P]ermanent markers placed at a disposal site, (2) public
16 records and archives, (3) government ownership and regulations regarding land or
17 resource use, and (4) other methods of preserving knowledge about the location, design,
18 and contents of a disposal system.” (40 CFR § 191.12)

19
20 **Pennsylvanian.** Second youngest period of the Paleozoic Era and also the corresponding
21 system of rocks. Pennsylvanian rocks are found about 12,800 to 15,000 feet below the
22 Los Medaños surface in the vicinity of the WIPP site and may contain oil and natural
23 gas. Subsequent to the Mississippian Period and followed by the Permian Period,
24 between approximately 320 and 286 million years ago.


25
26 **percolate.** The movement of liquid (usually water) through unsaturated or saturated solid and
27 rock.

28
29 **performance assessment.** An analysis that (1) identifies the processes and events that might
30 affect the disposal system; (2) examines the effects of these processes and events on the
31 performance of the disposal system; and (3) estimates the cumulative releases of
32 radionuclides, considering the associated uncertainties, caused by all significant
33 processes and events.

34
35 **performance modeling.** A process of building models of the factors affecting the
36 containment of nuclear waste to project into the future how the WIPP facility will
37 respond to probabilistic events and processes.

38
39 **permafrost.** Permanently frozen ground which occurs at different depths below the ground
40 surface in frigid climates.

41
42 **permeability.** In hydrology, the capacity of a rock, sediment, or soil to transmit fluids under
43 specified conditions. It is given by the hydraulic conductivity (K) times the fluid



1 dynamic viscosity (μ) divided by the fluid density (ρ) times acceleration due to gravity
2 (g); $k = \mu K / \rho g$.

3
4 **Permian Basin.** A region in the Central U.S. where, during Permian times 286 and 245
5 million years ago, there were many shallow seas that laid down vast beds of evaporites.
6 The Delaware basin is a part of the Permian basin.

7
8 **Permian Period.** This period is the last of seven periods of the Paleozoic Era, subsequent to
9 the Pennsylvanian Period; also, the corresponding system of rocks (between
10 approximately 286 and 245 million years ago).

11
12 **petrological.** Relating to the general study of rocks in all their aspects, including their
13 mineralogies, textures, and structures, their origins, field occurrences, alterations, and
14 their relationships to other rocks.

15
16 **pH.** The symbol used to designate a standard measure of the acidity of a solution. pH is
17 defined as the negative logarithm (base 10) of the activity of hydrogen ion in the
18 solution. Pure water, which is regarded as neutral, has a pH of 7. A lower value of pH
19 would indicate acidity, and a higher value would indicate that a solution is basic
20 (alkaline).

21
22 **phenols.** Organic compounds that are by-products of petroleum refining, tanning, and textile,
23 dye, and resin manufacturing. Low concentrations cause taste and odor problems in
24 water; higher concentrations can kill aquatic life and humans.

25
26 **physicochemical.** Describes chemically related processes studied for performance assessment
27 purposes that have potential to influence the transport of contaminants throughout the
28 disposal system. These physicochemical processes include alpha recoil, chemical
29 gradients and osmotic activity.

30
31 **physiography.** A description of the natural features of the surface of the earth.

32
33 **pisolitic.** Pertaining to pisolite [sed], or to the texture of a rock made up of pisoliths or
34 pealike grains; for example, pisolitic bauxite or pisolitic limestone.

35
36 **Pitzer Formalisms.** A semi-empirical formalism for calculating activity coefficients
37 particularly in saline aqueous systems.

38
39 **plasticity.** The property of a material that enables it to undergo permanent deformation
40 without appreciable volume change or elastic rebound, and without rupture.

41
42 **Pleistocene.** This is the first epoch of the Quaternary Period of the Upper-Cenozoic Era; also,
43 the corresponding system of rocks. It is subsequent to the Pliocene Epoch of the
44 Tertiary Period and precedes the Holocene Epoch of the Quaternary Period (between



Title 40 CFR Part 191 Compliance Certification Application

1 approximately 1.6 million to 10,000 years ago). Climatic changes during this period are
2 used as an analog in estimating future climatic extremes for the WIPP.

3
4 **Pliocene.** This is the last epoch of the Tertiary Period of the Middle-Cenozoic Era; also, the
5 corresponding system of rocks. It is subsequent to the Miocene Epoch of the Tertiary
6 Period and preceded by the Pleistocene Epoch of the Quaternary Period (between
7 approximately 5.3 and 1.6 million years ago).

8
9 **plug.** A watertight or gastight seal installed in a borehole or well to prevent movement of
10 fluids or gas. Also used to fill in or seal off fractures, cavities, or other porosity in the
11 walls of a borehole. Commonly a cement plug is used in the sealing of boreholes or
12 wells associated with oil, gas, or water exploration.

13
14 **plutonium.** A metallic, radioactive actinide, symbol Pu, atomic number 94, in the
15 transuranium series of elements; used as a nuclear fuel and as the fissile agent in nuclear
16 weapons.

17
18 **plutonium equivalent curie.** A term developed to provide a uniform basis among various
19 radioactive wastes to perform comparative human health consequence analyses resulting
20 from inhalation. The plutonium equivalent curie concept has strict limits of
21 applicability. It is utilized herein as a means of expressing the transuranic activity
22 content of transuranic waste packages.

23
24 **pmH.** The symbol used to designate a standard measure of the acidity of a solution. pmH is
25 defined as the negative logarithm of the molal concentration of hydrogen ion in solution.


26
27 **Poços de Caldas.** A location in Brazil where two uranium deposits were investigated for the
28 possibility that the flow of fluids might lead to the development and migration of a
29 large-scale redox front.

30
31 **point data.** Discrete actual data collected by directly observing and recording a measurable
32 event or attribute.

33
34 **point source.** A source of effluents that is small enough in dimensions that it can be treated
35 as if it were a point. The converse (not used in this document) is a diffuse source. A
36 point source can be either a continuous source or a source that emits effluents only in
37 pulses or for a short time.

38
39 **polyelectrolyte.** A substance of high molecular weight (as a protein or a nuclide).

40
41 **polyhalite.** An evaporite mineral: $K_2MgCa_2(SO_4)_4 \cdot 2H_2O$. It is a hard, nearly insoluble
42 mineral with no economic value.



Title 40 CFR Part 191 Compliance Certification Application

1 **population dose.** The sum of the radiation doses received by the individual members of a
2 population.

3
4 **porosity.** The ratio of the void volume in a rock or soil to its total volume. It may be stated
5 as a percentage or as a decimal.

6
7 **process.** A physical natural or anthropogenic phenomenon that occurs continuously or over a
8 significant portion of the time frame of interest, in other words, a “long-term”
9 phenomenon.

10
11 **portlandite.** An ingredient of Portland cement [Ca(OH)₂].

12
13 **possible futures.** The set of all possible occurrences within the 10,000 year regulatory time
14 frame.

15
16 **postclosure phase.** A designated period of time beginning with the end of the
17 Decommissioning Phase and extending through the end of the regulatory time frame of
18 10,000 years. Performance assessment modeling of repository behavior addresses this
19 time frame.

20
21 **potash.** A potassium compound used in agriculture and industry.

22
23 **potassium.** A silver-white metallic element of the alkali metal group. It occurs abundantly in
24 the McNutt Potash Zone in the vicinity of the WIPP site.

25
26 **potentiometric surface.** A subsurface map of the hydraulic potentials of an aquifer. It is
27 usually represented in figures as a contour map, each point estimating how high the
28 water would rise in a well tapping that aquifer at that point.

29
30 **Precambrian.** Applies to all geologic time, and its corresponding rocks, before the beginning
31 of the Paleozoic Era, 570 million years ago. The Precambrian basement is
32 approximately 18,000 feet (5,486 meters) below the surface in the vicinity of the WIPP
33 site.

34
35 **probabilistic analysis.** Analysis that arose from statistical investigations is referred to as
36 probabilistic analysis. This analysis is characterized by the fact that although specific
37 outcomes of an experiment cannot be predicted with certainty, relative frequencies for
38 various possible outcomes are predictable. Monte Carlo analysis is used for
39 probabilistic analysis of the WIPP.

40
41 **Project Gnome.** The Plowshare Project Gnome Test, an underground nuclear test, took
42 place on December 10, 1961 at a location approximately 8 miles (13 kilometers)
43 southwest of the WIPP site. Its primary purpose was to study the effects of an nuclear



Title 40 CFR Part 191 Compliance Certification Application

1 explosion in salt. The site was decommissioned in 1979, but continues to be monitored
2 by the EPA for radioactive contamination.

3
4 **promulgation.** Referring to the process of putting regulations into law.

5
6 **Public Law 96-164.** The U.S. Department of Energy National Security and Military
7 Applications of Nuclear Energy Act of 1980. Public Law 96-164 directed the DOE to
8 proceed with the design and development of the WIPP.

9
10 **Public Law 102-579.** See Land Withdrawal Act.

11
12 **quality assurance.** The planned and systematic actions necessary to provide adequate
13 confidence that a structure, system, or component will perform satisfactorily in service.

14
15 **Quality Assurance Project Plans.** This document identifies the quality of data necessary,
16 and the techniques designed to attain and ensure the required data quality objectives
17 associated with the WIPP waste characterization program.

18
19 **quality control.** Those quality assurance activities that provide a means to control and
20 measure the characteristics of a structure, system, or component to established
21 requirements.

22
23 **Quaternary.** The last period of the Cenozoic Era; also, the corresponding system of rocks. It
24 is subsequent to the Tertiary Period, beginning approximately 1.6 million years ago and
25 extends to the present.

26
27 **Rad.** See radiation absorbed dose.

28
29 **radially.** The term "radial" is the adjective of radius used to describe forces that act in
30 directions parallel to some radius or to designate lines, fractures, flow paths, etc., that
31 are oriented parallel to radii. "Radially" is the corresponding adverb. Effects that act
32 outward from a single point or inward toward a point (for example, flow of groundwater
33 toward a borehole) may be radial in form.

34
35 **radiation absorbed dose (Rad).** The special unit of absorbed dose of ionizing radiation.
36 One rad of absorbed dose is equal to the absorption of 100 ergs of radiation energy per
37 gram of absorbing material. See also rem.

38
39 **radioactive decay.** A process whereby the nucleus of an atom spontaneously emits excess
40 energy by emitting particles and/or rays (alpha, beta, gamma, or neutrons).

41
42 **radioactive material.** Matter composed of or containing radionuclides, with radiological
43 half-lives greater than 20 years, subject to the Atomic Energy Act of 1954, as amended.
44 (40 CFR § 191.12)



1 **radiographic examination.** The nondestructive technique that enables a qualitative
2 evaluation of the contents of a waste container.

3
4 **radiolysis.** Chemical decomposition by the action of radiation.

5
6 **radiometric.** Pertaining to the measurement of geologic time by the study of parent and/or
7 daughter isotopic abundances and known disintegration rates of the radioactive parent
8 isotopes, for example, *radiometric dating*.

9
10 **radionuclide.** An isotope having an unstable nucleus, that is subject to spontaneous decay.

11
12 **reactant.** A substance that enters into a chemical reaction and is altered. For example, metal
13 waste reacts with oxygen and water to produce hydrogen and metal oxides.

14
15 **real-time radiography.** A type of remote Non-Destructive Test that uses a luminescent
16 screen to form a moving image from x-rays that have passed through a rotating test
17 object. Real-time radiography is relied on for detection of waste form shapes and the
18 presence of liquids inside the waste package.

19
20 **reasonable.** (1) Not conflicting with reason, (2) not extreme or excessive, (3) having the
21 faculty of reason, or (4) possessing sound judgment.

22
23 **recharge.** The amount of water and the processes involved in the addition of water to an
24 aquifer.

25
26 **recharge point (or area).** In groundwater hydraulics, the point (or area) where surface water
27 enters an aquifer.

28
29 **recovery.** The process of getting something back. The term "resource recovery" is often used
30 to indicate the mining of minerals or the withdrawal of oil or natural gas from beneath
31 the surface of the Earth through wells.

32
33 **recrystallization.** The formation, which were initially in the solid state, of new crystalline
34 mineral grains in a rock. The new grains are generally larger than the original grains and
35 may have a different mineralogical composition. It is usually caused by increased
36 temperatures and pressures exerted on a rock formation.

37
38 **regulatory time frame.** As defined in 40 CFR Part 194, the regulatory time frame means the
39 time period beginning with disposal and ending 10,000 years after disposal.

40
41 **release.** Movement of regulated substances into the accessible environment as defined in 40
42 CFR Part 191 or beyond the unit boundary as defined for 40 CFR § 268.6.



1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43

rem. Roentgen equivalent in man—a special unit of dose equivalent which is the product of absorbed dose, a quality factor which rates the biological effectiveness of the radiation types producing the dose, and other modifying factors (usually equal to one). If the quality and modifying factors are unity, 1 rem is equal to 1 rad: 100 rem = 1 Sievert (SI units). Also expressed in terms of millirem (mrem): 1 rem = 1,000 mrem.

remote-handled transuranic waste. Packaged transuranic waste that has a measured external surface dose rate at or greater than 200 millirems per hour at the surface of the container.

repository. The portion of the WIPP underground system within the Salado Formation, including the access drifts, waste panels, and experimental areas, but excluding the shafts.

repository/shaft system. The WIPP underground workings, including the shafts, all engineered and natural barriers, and the altered zones within the Salado Formation and overlying units resulting from construction of the underground workings.

repressurization. The increase in pressure in a previously depleted volume, that might be associated with a fluid injection.

reserves. Mineral resources that can be extracted profitably by existing techniques and under present economic conditions.

reservoir. A subsurface volume of rock that has sufficient porosity and permeability to permit the accumulation of oil, gas, or water under adequate trap conditions. Also applies to any natural or artificial surface holding area used to store, regulate, or control water.

residuum. An accumulation of rock debris formed by weathering and remaining essentially in place after all but the least soluble constituents have been removed, usually forming a comparatively thin surface layer concealing the unweathered or partly altered rock below.

resistivity. Measure of electrical resistance in a fluid such as brine.

resources. Mineralization that is concentrated enough, in large enough quantity, and in a physical and chemical form such that its extraction is currently or potentially feasible and profitable.

retardation. Attenuation of contaminant concentration with migration distance. There are a variety of physical and chemical processes that can contribute to retardation.



1 **retrievable.** Describes storage of radioactive waste in a manner designed for potential
2 recovery without loss of control or release of radioactivity.

3
4 **Richter Scale.** A logarithmic scale expressing the magnitude of a seismic disturbance (as an
5 earthquake) in terms of the energy dissipated in it with 2 being the lowest and 8.5 being
6 the largest earthquakes yet recorded.

7
8 **Rio Grande Rift.** The rift system of the Rio Grande valley in which the crustal plate has split
9 and a series of grabens have formed in New Mexico and Southern Colorado. It is a
10 product of extensional tectonics.

11
12 **room.** An excavated cavity within a panel in the underground. Within the Waste Isolation
13 Pilot Plant, a room is about 10 meters wide, 4 meters high, and 91 meters long.

14
15 **rubidium-strontium age.** Determination of an age for a mineral or rock in years based on the
16 ratio of radiogenic strontium-87 to rubidium-87 and the known radioactive decay rate of
17 rubidium-87. If ratios are measured for more than one phase of a single rock, or for a
18 number of related rocks that differ in rubidium content, an isochron may be drawn.

19
20 **Rustler Formation.** The evaporite beds, including mudstones, of probable Permian age that
21 immediately overlie the Salado Formation.

22
23 **Salado Formation.** A geologic formation of Late Permian age in southeastern New Mexico.
24 At the WIPP site, it is composed of salt beds with minor amounts of anhydrite (45
25 numbered anhydrite marker beds: Marker Bed 101 through Marker Bed 145) and clay.
26 It is the host unit for the Waste Isolation Pilot Plant repository.

27
28 **saline.** Waters are considered saline when they contain between 10 and 100 parts per
29 thousand salinity. See brackish and brine.

30
31 **saline intrusion.** A human-induced coastal process whereby subsurface seawater
32 progressively replaces freshwater withdrawn from an aquifer being pumped close to the
33 shoreline.

34
35 **Sandia National Laboratories.** A multiprogram laboratory located in Albuquerque, NM,
36 and Livermore, CA. SNL is operated and managed for the DOE by the Sandia
37 Corporation. From 1949 until October 1993, Sandia Corporation was a wholly owned
38 subsidiary of AT&T. Sandia Corporation is currently a wholly owned subsidiary of
39 Lockheed-Martin Corp.

40
41 **San Simon Sink.** The central, most depressed area of San Simon Swale.

42
43 **San Simon Swale.** A broad depression about 15 miles east of the Los Medaños site, open to
44 the southeast.



1 **Santa Rosa Formation.** The Santa Rosa Sandstone is a principal aquifer in several areas,
2 particularly in Winkler and Ward Counties, Texas. It produces both fresh and saline
3 water, depending on location. The westernmost extent of the Santa Rosa Sandstone is
4 just into Eddy County. Wells completed in the Santa Rosa Sandstone have low yields
5 with specific capacities of 0.14-0.2 gpm per foot of drawdown; the formation porosity is
6 about 13 percent.

7
8 **saturated.** A condition in which all connected pores in a given volume of material contain
9 fluid.

10
11 **scarp.** A line of cliffs produced by faulting or by erosion. The term is an abbreviated form of
12 escarpment, and the two terms commonly have the same meaning, although "scarp" is
13 more often applied to cliffs formed by faulting.

14
15 **scenario.** A combination of naturally occurring or human-induced events and processes that
16 represent realistic future changes to the repository, geologic, and geohydrologic systems
17 that could cause or promote the escape of radionuclides and/or hazardous constituents
18 from the repository.

19
20 **screening argument.** Regulatory Guidance, Probability of Occurrence, and Consequence, are
21 three criteria used to eliminate from scenario development those events and processes
22 that are not applicable to a specific disposal system or that do not have the potential of
23 contributing significantly to the performance measure (for example, integrated
24 radionuclide releases).

25
26 **seal.** An engineered barrier designed to isolate the waste and to impede groundwater flow in
27 the shafts.


28
29 **sealing system.** Includes all components of the WIPP seal design program.

30
31 **sedimentary.** A descriptive term for rocks formed of sediment such as clastic rocks
32 (sandstones and shales), secretions of organisms, limestones, and precipitation from
33 solution, rock salts and gypsums.

34
35 **sedimentation.** The process of accumulating sediment into layers of solid rock which
36 includes the generation of rock particles, the transportation and the deposition of these
37 particles, and the consolidation of these particles into solid rock.

38
39 **seismic.** Applies to the activity of naturally or artificially induced earthquakes or earth
40 vibrations, where the seismic waves are the elastic waves produced by these vibration.

41
42 **seismic risk zone.** A designation of a geographic region expressing the maximum intensity of
43 earthquakes that could be expected there.



Title 40 CFR Part 191 Compliance Certification Application

1 **sensitivity analysis.** Methods for computing the effect of changes in the input parameters on
2 the model predictions.

3
4 **seral.** Relating to a series of ecological communities formed in ecological succession.

5
6 **sequestration.** To hold (as a metallic ion) in solution.

7
8 **shaft.** A man-made hole, either vertical or steeply inclined, that connects the surface with the
9 underground workings of a mine.

10
11 **shaft pillar.** The cylindrical volume of rock around a shaft from which major underground
12 openings are excluded in order that they do not weaken the shaft.

13
14 **shallow-dissolution zone.** A zone of residual material at the interface of the Rustler and
15 Salado formations left after dissolution of salt. It is highly permeable and often contains
16 brine.

17
18 **SI unit.** See definition for System International Unit of Measure.

19
20 **Sievert.** The SI unit of effective dose. It is equal to 100 rem or one joule per kilogram. (40
21 CFR § 191.12)

22
23 **siliciclastic.** Pertaining to clastic noncarbonate rocks which are almost exclusively silicon-
24 bearing, either as forms of quartz or as silicates.

25
26 **siltstone.** An indurated silt having the texture and composition of shale but lacking its fine
27 lamination or fissility; a massive mudstone in which the silt predominates over clay; a
28 nonfissile silt shale.

29
30 **Silurian.** Third oldest period of the Paleozoic Era, and also the corresponding system of
31 rocks. Subsequent to the Ordovician Period and followed by the Devonian Period,
32 between approximately 438 and 400 million years ago. The Silurian of the southeast
33 New Mexico - west Texas area consists of the Fusselman limestone and the carbonates
34 and shales of an Upper Silurian unit, both of which were deposited in the broad
35 subsiding area named the Tobosa Basin.

36
37 **Simpson.** In the Delaware basin, the Simpson consists of three main layers of limestone,
38 alternating with thinner green, brown and black shale, black shale with rounded quartz
39 grain inclusions, and sandstone.

40
41 **site characterization.** The process of making geologic and environmental studies to identify
42 potential sites for mined geologic repositories. Detailed site characterization goes
43 further: additional data are collected that would be necessary if a license application
44 were to be submitted.



1 **SKI (Statens Kärnkraftinspektion).** An abbreviation for the Swedish Nuclear Power
2 Inspectorate, Statens Kärnkraftinspektion. A list of potentially relevant features, events,
3 and processes developed by SKI was considered to be the best documented and most
4 comprehensive starting point for the WIPP scenario development process.
5

6 **sludge.** Refers to de-watered contact-handled transuranic wastes containing both organic and
7 inorganic constituents that must meet the Waste Acceptance Criteria for shipment and
8 disposal at the WIPP repository. High sludges are contact-handled transuranic waste
9 where the sludge component constitutes 50 percent or more of the waste volume; low
10 sludges are the same type of waste containing less than 50 percent by volume of sludge.
11

12 **SO-C.** Screened-Out on the basis of Consequence. The term applies to the elimination of
13 features, events, and processes from performance assessment and compliance
14 assessment calculations for WIPP compliance efforts on the basis of consequence.
15

16 **solifluction.** The slow downslope flow of waterlogged soil that is commonly active in
17 periglacial regions underlain by frozen ground that acts as a downward barrier to water
18 percolation.
19

20 **SO-P.** Screened-Out on the basis of low Probability. The term applies to the elimination of
21 features, events, and processes from performance assessment and compliance
22 assessment calculations for WIPP compliance efforts on the basis of low probability of
23 occurrence.
24

25 **SO-R.** Screened-Out on the basis of Regulations. The term applies to the elimination of
26 features, events, and processes from performance assessment calculations used for WIPP
27 compliance efforts on the basis of regulations provided in 40 CFR Part 191 and criteria
28 provided in 40 CFR Part 194.
29

30 **solubility.** The ability or tendency of one substance to blend uniformly with another (for
31 example, solid in liquids, liquid in liquid, gas in liquid, and gas in gas). Solids vary
32 from 0 to 100 percent in their degree of solubility in liquids depending on the chemical
33 nature of the substance(s); to the extent that they are soluble, they lose their crystalline
34 form and become molecularly or ionically dispersed in the solvent to form a true
35 solution. Liquids and gases are often said to be miscible in other liquids and gases
36 rather than soluble.
37

38 **solute.** A substance which is dissolved in another substance called the solvent. The solute is
39 uniformly dispersed in the solvent either molecularly or ionically.
40

41 **solution.** A mixture, liquid, solid, or gaseous, in which the components are uniformly
42 distributed throughout the mixture.
43



1 **solution subsidence.** Gradual subsidence of nonsoluble strata due to the dissolution of
2 underlying rock.

3
4 **solvent.** A substance capable of dissolving another substance (solute) to form a uniform
5 dispersed mixture (solution) at the molecular or ionic level. Solvents are, accordingly,
6 characterized as either polar or non-polar. Water is strongly polar; hydrocarbon solvents
7 are non-polar.

8
9 **soret effect.** The principle that asserts when temperature differences are induced in a solution
10 of common salt or other substance in water, the dissolved material will become
11 relatively concentrated in those portions in which the temperature is lowest.

12
13 **sorption.** The binding on a microscopic scale of one substance to another, such as by
14 adsorption or ion exchange. In this document, the word is especially used in the
15 sorption of solutes onto aquifer solids.

16
17 **source term.** The kinds and amounts of radionuclides that make up the source of a potential
18 release of radioactivity. *See* Nuclide inventory.

19
20 **spallings.** During exploratory drilling, waste surrounding the eroded borehole that is
21 transported by waste-generated gas escaping to the lower pressure borehole.

22
23 **specific activity.** Radioactivity per unit weight of radioactive material.

24
25 **standard waste box.** A waste container measuring approximately 6 by 4.5 by 3 feet high,
26 with rounded ends.

27
28 **steady-state.** A state or condition of a system or process that does not change in time; a
29 condition that changes only negligibly over a specified time.

30
31 **stochastic uncertainty.** As used in probabilistic risk assessment, stochastic uncertainty
32 characterizes uncertainty involving conceptual models and/or the future.

33
34 **strata.** Geologic term for layers of the earth's crust. The crust was generally laid down in
35 layers during geological epochs.

36
37 **stratigraphy.** The science and study of the origin, composition, and proper sequence in
38 which various rock strata were layered during various geological ages.

39
40 **strike slip fault.** A fault on which the movement is parallel to the fault's strike.

41
42 **structure map.** A map that portrays subsurface configuration by means of structure contour
43 lines.



1 **subduction zone.** A long, narrow belt in which subduction takes place, for example, along
2 the Peru-Chile trench, where the Pacific plate descends the South American plate.

3
4 **study area.** The region about the Los Medaños site studied in the evaluation of that site.

5
6 **subjective uncertainty.** Subjective uncertainty derives from a lack of knowledge about
7 quantities, attributes, or properties believed to have a single or certain range of values.

8
9 **subsidence.** The downward settling of the earth's surface with little or no horizontal motion.
10 Subsidence may be caused by natural geological processes or by man-made activities.

11
12 **substrate.** The surface and/or nutrient on which an organism will live and grow.

13
14 **subsurface.** The rock and soil below the ground surface whose geological features,
15 principally stratigraphy and structure, are interpreted primarily on the basis of drilling
16 activities and geophysical evidence.

17
18 **subvertical.** Used to describe planar features oriented within approximately 15 degrees of
19 vertical.

20
21 **sulfates.** A mineral compounds characterized by the sulfate radical SO_4 . Anhydrous sulfates,
22 such as barite, $BaSO_4$, have divalent cations linked to the sulfate radical; hydrous and
23 basic sulfates, such as gypsum, $CaSO_4 \cdot 2H_2O$, contains water molecules.

24
25 **supersaturation.** A solution, such as the Salado Formation brine, that contains more of a
26 solute (radioactive isotope) than is normally present when equilibrium is established.

27
28 **swipe samples.** The presence of radioactive contaminants may be ascertained by applying a
29 Kim-wipe™ or equivalent to the surface of the potentially contaminated item and
30 measuring the radioactivity of the Kim-wipe™.

31
32 **sylvite.** A mineral, KCl, used in the manufacture of fertilizer.

33
34 **syndeposition.** Processes occurring during deposition of a sedimentary rock, and contributing
35 to the formation of the sedimentary framework.

36
37 **System of International units.** The version of the metric system which has been established
38 by the International Bureau of Weights and Measures and is administered in the United
39 States by the National Institute of Standards and Technology. The abbreviation for this
40 system is "SI." (40 CFR § 191.12)

41
42 **Tamarisk Member.** Middle anhydrite layer of Rustler Formation.



1 **Tansill Formation.** The Tansill Formation is the upper most formation of the Artesia Group,
2 developing in the Mid-Permian Period, between approximately 280 and 260 million
3 years ago. The Artesia Group represents the shelf region of the Goat Seep Formation
4 and the Capitan Limestone. To the south of the Delaware Basin the Tansill Formation
5 consists of dolomites and sandstones. Toward the northern part of the basin, gypsum
6 and anhydrite along with siltstones and dolomites increase in the Tansill Formation. The
7 Tansill Formation is time correlative to the Capitan Reef at the basin margin and the
8 Bell Canyon Formation in the Delaware Basin.
9

10 **tectonics.** The branch of geology dealing with regional structural or deformational features.
11

12 **tensile.** Refers to tension in the mechanical sense. A material is under tension when the
13 stress exerted on it (tensile stress) acts “outward” such that the effect is to increase the
14 length or volume of the body.
15

16 **Tertiary Period.** This period is the first of two periods in the Cenozoic Era and also, the
17 corresponding system of rocks. It is subsequent to the Cretaceous Period of the
18 Mesozoic Era and is followed by the Quaternary Period of the Cenozoic Era. The
19 Tertiary Period dates between approximately 66.4 and 1.6 million years ago.
20

21 **thermal effect.** The effects of naturally occurring heat on groundwater flow.
22

23 **thermal field.** The field or set of temperatures throughout a volume. Use of the term usually
24 connotes temperatures that differ from point to point.
25

26 **thermal gradient.** The rate of change of temperature in the direction of increasing
27 temperature.
28

29 **thermodynamic.** A term that deals with the relations between heat and other forms of
30 energy. A system is at thermodynamic equilibrium when there is no tendency for
31 change in the free energy of the system,
32

33 **thief zone.** A rock unit responsible for excessive fluid loss from boreholes during drilling or
34 production operations.
35

36 **topographic/topography.** Relating to, or concerning the configuration of a surface that
37 includes relief and the position of its natural and man-made features.
38

39 **transient.** This term is used to describe changing state or condition as in ground motion due
40 to seismic activity, or groundwater flow resulting from changing fluid densities.
41

42 **transmissivity.** The rate at which water is transmitted through a unit thickness of an aquifer.
43 It is considered a property of the aquifer which is transmissive. The contained liquid is
44 transmissible.



1 **transport.** For regulatory purposes relating to moving waste to the repository. The
2 movement of radioactive wastes across public routes via the use of packaging as the
3 transportation container and the movement of a hazardous substance by any mode,
4 including pipeline (as defined in the Pipeline Safety Act). In the case of a hazardous
5 substance that has been accepted for transportation by a common or contract carrier, the
6 term "transport" or "transportation" shall include any stoppage in transit that is
7 temporary, incidental to the transportation movement, and at the ordinary operating
8 convenience of a common or contract carrier, and any such stoppage shall be considered
9 as a continuity of movement and not as the storage of a hazardous substance. When
10 addressing the performance of the geologic repository, "transport" refers to the
11 movement of material by mechanisms such as solution, or association with colloids, in
12 groundwater or water from other sources flowing through the repository, seepage of gas
13 through fractured or porous media, or entrainment in drilling fluid, brine, or gas flowing
14 in a borehole.

15
16 **transuranic.** See Transuranic Waste.

17
18 **transuranic nuclide.** A nuclide with an atomic number greater than that of uranium (92).
19 All transuranic nuclides are produced artificially and are radioactive.

20
21 **Transuranic Package Transporter-II.** Package designed to transport contact-handled
22 transuranic mixed waste to the WIPP site. It is a cylinder with a flat bottom and a
23 domed top that is transported in the upright position. Also called TRUPACT-II.

24
25 **transuranic waste.** The term "transuranic waste" means waste containing more than
26 100 nanocuries of alpha-emitting transuranic isotopes per gram of waste, with half-lives
27 greater than 20 years, except for: (1) high-level radioactive waste, (2) waste that the
28 Secretary has determined, with the concurrence of the Administrator, does not need the
29 degree of isolation required by the disposal regulations, or (3) waste that the Nuclear
30 Regulatory Commission has approved for disposal on a case-by-case basis in accordance
31 with 10 CFR 61.

32
33 **treatment.** Means any method, technique, or process, including neutralization, designed to
34 change the physical, chemical, or biological character or composition of any hazardous
35 waste so as to neutralize such waste, or so as to recover energy or material resources
36 from the waste, or so as to render such waste non-hazardous, or less hazardous; safe to
37 transport, store, or dispose of; or amenable for recovery, amenable for storage, or
38 reduced in volume.

39
40 **Triassic.** The first period of the Mesozoic era (after the Permian of the Paleozoic Era, and
41 before the Jurassic), thought to have covered the span of time between 225 and 190
42 million years ago; also, the corresponding system of rocks. The triassic is so named
43 because of its threefold division in the rocks of Germany.



1 **true solution.** A uniformly dispersed mixture at the molecular or ionic level, of one or more
2 substances (solute) in one or more substances (solvent). Solutions that exhibit no
3 change of internal energy upon mixing and have complete uniformity of cohesive forces
4 are true.

5
6 **Type A Packaging.** Means a packaging designed to retain the integrity of containment and
7 shielding required by this part under normal conditions of transport as demonstrated by
8 the tests set forth in 49 CFR § 173.465 or 173.466, as appropriate.

9
10 **uncertainty analysis.** (1) An evaluation to determine the uncertainty in model predictions
11 that results from imprecisely known input variables. (2) Determination of the degree of
12 uncertainty in the results of a calculation based on uncertainties in the input parameters
13 and underlying assumptions. Such an analysis requires definition of a system,
14 description of the uncertainties in the factors that are to be investigated, and the
15 characteristics of the system that are to be simulated.

16
17 **underpressurized.** A zone or rock formation (Culebra Dolomite or perhaps units underlying
18 the Salado Formation) that falls below the expected pressure gradient as depth
19 increases.

20
21 **undisturbed performance.** “[T]he predicted behavior of a disposal system, including
22 consideration of the uncertainties in predicted behavior, if the disposal system is not
23 disrupted by human intrusion or the occurrence of unlikely natural events.” (40 CFR §
24 191.12)

25
26 **Uninterruptible Power Supply.** A power supply that provides automatic, instantaneous
27 power, without delay or transients, on failure of normal power. It can consist of
28 batteries or full-time operating generators. It can be designated as standby or emergency
29 power depending on the application. Emergency installations must meet the
30 requirements specified for emergency.

31
32 **unit boundary.** In the context of 40 CFR § 268.6, the unit boundary is that point at which
33 migration occurs if hazardous constituents pass that point in concentrations exceeding
34 health-based levels.

35
36 **uplift.** A structurally high area in the crust, produced by positive movements that raise or
37 upthrust the rocks, as in a dome or arch.

38
39 **Upper Devonian Woodford Shale.** The Woodford is a dark brown to black, fissile,
40 bituminous, spore-bearing shale which becomes [arenaceous] northward and contains
41 black [chert] to the south and west.

42
43 **U.S. Department of Energy.** The cabinet-level U.S. Government agency responsible for
44 weapons production, energy research, and for the subsequent cleanup of hazardous and



1 radioactive waste sites. It was created from the Energy Research and Development
2 Administration and other federal government functions in 1977. DOE is also
3 responsible for the construction and oversight of WIPP primarily through its Carlsbad
4 Area Office.

5
6 **U.S. Environmental Protection Agency.** (1) An independent agency of the Federal
7 government formed in 1970 by Presidential executive order, bringing together parts of
8 various government agencies involved with the control of pollution, and responsible for
9 pollution abatement and control programs, including programs in air and water pollution
10 control, water supply and radiation protection, solid and toxic waste management,
11 pesticides control, and noise abatement; (2) The Agency for those implementation
12 responsibilities for the Waste Isolation Pilot Plant, given to the Agency by the Waste
13 Isolation Pilot Plant Land Withdrawal Act (Pub. L. 102-579, 106, Stat. 4777) which are:

14
15 *(i) Determinations by the Agency that the Waste Isolation Pilot Plant is in compliance with*
16 *Subpart A of this part;*

17
18 *(ii) Issuance of criteria for the certifications of compliance with subparts B and C of this part*
19 *of the Waste Isolation Pilot Plant's compliance with subparts B and C of this part;*

20
21 *(iii) Certifications of compliance with subparts B and C of this part of the Waste Isolation*
22 *Pilot Plant's compliance with subparts B and C of this part;*

23
24 *(iv) If the initial certification is made, periodic recertification of the Waste Isolation Pilot*
25 *Plant's continued compliance with subparts B and C of this part;*

26
27 *(v) Review and comment on performance assessment reports of the Waste Isolation Pilot*
28 *Plant; and,*

29
30 *(vi) Concurrence by the Agency with the Department's determination under § 191.02(i) that*
31 *certain wastes do not need the degree of isolation required by subparts B and C.*

32
33 **U.S. Energy Research and Development Agency.** Energy Research and Development
34 Agency, a forerunner of the DOE that was formed in 1974 (Pub. L. 93-438).

35
36 **U.S. Geological Survey.** U.S. Geological Survey, Department of Interior.

37
38 **valence state.** The degree of oxidation (oxidation state) of an element or atom. It is usually
39 expressed as a positive or negative number representing the ionic charge or effective
40 charge of the element of atom.

41
42 **Van Horn Sandstone.** Clastics of the late Precambrian age that crop out near Van Horn,
43 Texas and are overlain by the Bliss Sandstone.



1 **varve.** In its broadest sense, a varve is a usually thin layer of sedimentary rock that was
2 deposited in a single year. The term is most commonly used to describe sediments
3 deposited in glacial meltwater lakes that exhibit a sequence of alternating coarser and
4 finer layers representing summer and winter deposition, respectively.
5

6 **vector.** A quantity or guide that requires a magnitude, direction and sense. It is commonly
7 represented by a line, but also includes the direction of a substance (or activity) such as
8 the flow of water in a general direction. As used in performance assessment, the term is
9 also used to describe model results from one set of parameters selected through
10 statistical techniques.
11

12 **viscosity.** A property of fluids which is displayed by definite resistance to change of form and
13 many solids show a gradual yielding to forces tending to change their form. This
14 property, a sort of internal friction, is called viscosity.
15

16 **volatile organic compounds.** An organic compound that evaporates (volatilizes) readily at
17 room temperature.
18

19 **vug.** Small open cavity in a rock.
20

21 **Waste Acceptance Criteria.** A set of conditions established for permitting transuranic
22 wastes to be packaged, shipped, managed, and disposed of at the WIPP.
23

24 **waste characterization.** Sampling, monitoring, and analysis activities to determine the
25 nature of the waste to be emplaced at the WIPP.
26

27 **waste disposal panels.** A collection of disposal rooms in the WIPP repository. Each panel
28 will be sealed as a unit.
29

30 **waste form.** A term used to emphasize the physical and chemical properties of the waste.
31

32 **Waste Isolation Pilot Plant.** (1) The project authorized under § 213 of the DOE National
33 Security and Military Application of Nuclear Energy Authorizations Act of 1980 (Public
34 Law 96-164; Stat. 1259, 1265) to demonstrate the safe disposal of radioactive waste
35 materials generated by atomic defense activities; (2) A research and development
36 facility, located near Carlsbad, New Mexico, to be used for demonstrating the safe
37 disposal of transuranic wastes from DOE activities.
38

39 **waste matrix.** The material that surrounds and contains the waste and to some extent protects
40 it from being released into the surrounding rock and groundwater. Only material within
41 the canister (or drum or box) that contains the waste is considered part of the waste
42 matrix.
43



1 **waste stream.** A grouping of waste materials with specific definable characteristics that
2 remain the same throughout the process generating the waste stream. A waste stream is
3 produced by a single process or subprocess or may be combined to produce a single
4 output waste stream.

5
6 **waste unit factor.** The waste unit factor is the normalizing factor used to determine the
7 release limits for 40 CFR § 191.13(a). The term is the same as the “unit of waste”
8 defined in 40 CFR Part 191 Appendix B, Table 1, note e, as “an amount of transuranic
9 wastes containing one million curies of alpha-emitting transuranic radionuclides with
10 half-lives greater than 20 years.” For the WIPP, the overall quantity of radionuclides
11 that apply to the waste unit is 4.07×10^6 curies, and the waste unit factor is therefore
12 4.07.

13
14 **wicking.** The mechanism by which a liquid can migrate through an unsaturated zone against
15 the forces of gravity and surface tension in the soil. Also called capillary rise.

16
17 **Wink Sink.** Wink Sink is located 100 miles (160 kilometers) southeast of the WIPP site
18 outside the Delaware Basin. The surface impression is 360 feet (110 meters) in width
19 and 110 feet (34 meters) deep. It is the result of dissolution of the Salado Formation
20 caused by percolation of shallow groundwater through abandoned boreholes.
21 Subsequent subsidence of the rock units overlying the Salado Formation led to the
22 collapse feature present on the surface.

23
24 **WIPP site boundary.** The boundary that defines the outer limits of the 16-section land
25 withdrawal area. For the purposes of this application, this boundary defines the
26 controlled area.

27
28 **Wolfcampian.** Lower member of Permian age in Southeastern New Mexico.

29
30 **working agreement.** Appendix B of the Agreement of Consultation and Cooperation, which
31 sets forth the working details of that Agreement.

32
33 **X-ray.** Any of the electromagnetic radiations of the same nature as visible radiation but of
34 extremely short wavelength (less than 100 angstroms). X-rays are produced by
35 bombarding a metallic target with fast electrons in vacuum or by transition of atoms to
36 lower energy states, and have the properties of ionizing a gas upon passage through it, of
37 penetration various thicknesses of all solids, of producing secondary radiations by
38 impinging on materials bodies, of acting on photographic films and plates as light does,
39 and of causing fluorescent screens to emit light.

40
41 **Yates Formation.** The Yates Formation is one of five formations in the Artesia Group
42 developing in the Mid-Permian Period, between approximately 280 and 260 million
43 years ago. It is subsequent to the Seven Rivers Formation and is preceded by the Tansill
44 Formation. The Tansill Formation, Seven Rivers Formation, and the Yates Formation

1
2
3
4
5
6
7

represent the shelf region of the Capitan Limestone at the margin of the Delaware Basin. To the south it consists of dolomites and sandstones. Siltstones, mudstones, and dolomites increase toward the northern part of the basin. The Yates Formation is time correlative to the Capitan Reef at the basin margin and the Bell Canyon Formation in the Delaware Basin.



1
2

THIS PAGE INTENTIONALLY LEFT BLANK

